



EPA REG. NO. 73806-1

## FIFRA 2(ee) RECOMMENDATION FOR USE IN CALIFORNIA

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the U.S. Environmental Protection Agency.

**Citrus Crops (including subtypes, hybrids and citrus-like crops), Avocado and Other Tropical and Subtropical Fruits (Bearing and Non-Bearing).**

**Grapes and Cane Berry Crop Grouping:** Grapes (all types); Blackberry (including bingberry, black satin berry, boysenberry, Cherokee blackberry, Chesterberry, Cheyenne blackberry, coryberry, darrowberry, dewberry, Dirksen thornless berry, Himalayaberry, hullberry, Lavacaberry, lowberry, Lucretiaberry, mammoth blackberry, marionberry, nectarberry, olallieberry, Oregon evergreen berry, phenomenalberry, rangeberry, ravenberry, rossberry, Shawnee blackberry, youngberry, and varieties and/or hybrids of these), Blueberry, Cane Berries, Cranberry, Currant, Elderberry, Gooseberry, Huckleberry, Loganberry and Raspberry (black and red).

**Pome Fruits Crop Grouping (Bearing and Non-Bearing)** including Apple, Crabapple, Loquat, Mayhaw, Papaya, Pear, Pomegranate and Quince.

**Stone Fruits Crop Grouping** including Olives: Apricot, Cherry, Nectarine, Olive, Peach, Plum, Pluot and Prune.

### DIRECTIONS FOR USE

- It is a violation of Federal law to use this product inconsistent with its labeling.
- This recommendation must be in the possession of the user at the time of pesticide application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered labels.

### GENERAL APPLICATION INSTRUCTIONS

CROP	DISEASE	APPLICATION METHOD	RATE OF APPLICATION	APPLICATION PROGRAM
Grapes and Caneberries Grouping; Pome Fruits Crop Grouping; Stone Fruits Crops Grouping.	Alternaria, Anthracnose, Botrytis cinerea Botryosphaeria, (including dothidea) Pseudomonas spp., (including syringae), Rust, Scab	Foliar	1 – 3 quarts in a minimum of 100 gallons of water per acre. Low volume use per acre rate of K-PHITE 7LP and adjust amount of water to equipment needs.	Apply at 2-3 week intervals on a treated acre basis, use lower rates on first and second leaf trees, use higher rates on third leaf and greater. Do not apply K-PHITE 7LP at intervals of less than three days.
		Aerial	1-3 quarts in a minimum of 10 gallons of water/acre	Apply at 2-3 week intervals. Do not apply K-PHITE 7LP at intervals less than 3 days.
		Irrigation	Overhead: 2-4 quarts in at least 100 gallons of water. Drip: 2-3 quarts in at least 100 gallons of water.	Apply at 2-3 week intervals on a treated acre basis, use lower rates on first and second leaf trees, use higher rates on third leaf and greater. Do not apply K-PHITE 7LP at intervals of less than three days.
*Make applications prior to disease development in conjunction with good cultural management practices. *Use the higher rate when disease pressure is severe.				

All applicable directions, restrictions, and precautions on the EPA-registered label are to be followed. This FIFRA Section 2(ee) recommendation contains new or additional Diseases, which may not appear on the package label. Read and observe the cautionary statements plus all other information appearing on the product label.

REGISTRANT : **PLANT FOOD SYTEMS, INC**

P.O. BOX 775, 2827 Union Street, Zellwood, FL. 32798

