



EPA REG. NO. 73806-1

FIFRA 2(ee) RECOMMENDATION FOR USE IN CALIFORNIA

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the U.S. Environmental Protection Agency.

Cucurbit Crop Grouping: Cantaloupe, Chayote, Chinese waxgourd, Citron melon, Cucumber (all types), Gherkin, Gourd, Honeydew, Muskmelon, Pumpkin, Squash (summer and winter) and Watermelon.

APPLICATION GUIDELINES: Disease Prevention Program: Apply lower rate at 7-28 day intervals. Do not apply K-PHITE 7LP at intervals less than 3 days. Disease Control Program: Apply higher rate at 7-14 day intervals until control is reached. Under severe circumstances, application can be made at intervals of up to every three days. Consult with Farm Advisor, Licensed Agricultural Pest Control Advisor or Certified Crop Advisor to determine disease severity. Do not apply K-PHITE 7LP at intervals less than 3 days.

Make applications prior to disease development in conjunction with good cultural management practices. Use the higher rate when disease pressure is severe. Use rates and frequency are designed to prevent plant injury. If more frequent applications are necessary for disease control, consult a crop expert and follow their recommendations. Do not exceed the highest rate per application. Do not apply at less than 3 day intervals. Do not apply to plants that are dormant or heat or moisture stressed. Allow foliage to completely dry after application. Do not apply when conditions favor wet tissue for prolonged periods (>4 hours).

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product inconsistent with its labeling.
- This recommendation must be in the possession of the user at the time of pesticide application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered labels.

GENERAL APPLICATION INSTRUCTIONS

DISEASE	APPLICATION METHOD	RATE OF APPLICATION
Fusarium Phytophthora <i>Pseudomonas</i> spp. (including <i>syringae</i>)	Chemigation	2-4 quarts in at least 400 gallons of water/acre.
Pythium Ralstonia solanacearum Rhizoctonia Sclerotinia	Irrigation	Overhead: 2-4 quarts in at least 100 gallons of water. Drip: 2-3 quarts in at least 100 gallons of water.
Xanthomonas spp. (including <i>campestris</i>)	Planter Bed, Root Dip or Transplant Drench	1-2 ounces /1000 ft. of row. 1-2 quarts/100 gallons of water, 1/3 – 2/3 fluid ounce/gallon of water. Can be applied prior to, during or after transplanting, in seed furrow, knife injected near seed, or over covered seed bed.
	Soil Directed or Injected Application	2-4 quarts in min 20 gallons/acre. Lightly irrigate after application.

All applicable directions, restrictions, and precautions on the EPA-registered label are to be followed. This FIFRA Section 2(ee) recommendation contains new or additional Diseases, which may not appear on the package label. Read and observe the cautionary statements plus all other information appearing on the product label.

REGISTRANT : PLANT FOOD SYTEMS, INC

P.O. BOX 775, 2827 Union Street, Zellwood, FL. 32798

