

# SUPPLEMENTAL LABELING

ACEQUINOCYL | GROUP 20B | INSECTICIDE

## KANEMITE® 15 SC MITICIDE

Supplemental Label to Add DIRECTIONS FOR USE on GUAVA and TROPICAL and SUBTROPICAL, SMALL FRUIT, INEDIBLE PEEL (Crop Subgroup 24A)

This supplemental label expires June 22, 2021 and must not be used or distributed after this date.

<b>ACTIVE INGREDIENT:</b>	<b>% BY WT.</b>
*Acequinocyl .....	15.8%
<b>OTHER INGREDIENTS:</b> .....	84.2%
<b>TOTAL:</b> .....	100.0%

\*3-dodecyl-1, 4-dihydro-1, 4-dioxo-2-naphthyl-acetate.  
Contains 1.25 lb ai/gal.

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION / PRECAUCIÓN**

**DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

THIS LABELING MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF APPLICATION. READ THE LABEL AFFIXED TO THE CONTAINER FOR **KANEMITE® 15 SC MITICIDE** BEFORE APPLYING. USE OF **KANEMITE 15 SC MITICIDE** ACCORDING TO THIS LABELING IS SUBJECT TO THE USE PRECAUTIONS AND LIMITATIONS IMPOSED BY THE LABEL AFFIXED TO THE CONTAINER FOR **KANEMITE 15 SC MITICIDE (EPA Reg. No. 66330-38)**.

**061818V004SuFL**

**GUAVA (including: CATTLEY GUAVA, PARA GUAVA, PURPLE STRAWBERRY GUAVA, STRAWBERRY GUAVA AND YELLOW STRAWBERRY GUAVA).**

**TROPICAL AND SUBTROPICAL, SMALL FRUIT, INEDIBLE PEEL, CROP SUBGROUP 24A: Aisen, Bael fruit, Burmese grape, Cat's eyes, Inga, Longan, Lychee, Madras-thorn, Manduro, Matisia, Mesquite, Mongongo fruit, Small-flower pawpaw, Satinleaf, Sierra Leone-tamarind, Spanish lime, Velvet tamarind, Wampi, White star apple and cultivars, varieties and hybrids of these commodities).**

PEST	USE RATE
False spider mite ( <i>Brevipalpus spp.</i> )	31 fl oz/A (0.3 lb ai/A)

**RESTRICTIONS**

- Do not use less than 100 gallons of water volume per acre.
- Allow a minimum of 21 days between treatments.
- Do not apply this product through any type of irrigation system.
- Do not apply by air.
- Do not apply within 75 feet of aquatic areas.
- Not for indoor residential use.
- A 30-day plant-back interval is required for all crops not listed on the label.
- Do not make more than two applications of **KANEMITE 15 SC MITICIDE** per year.
- Do not apply more than 62 fl oz of **KANEMITE 15 SC MITICIDE** per acre (0.6 lb ai) per season.
- Preharvest interval is 1 day.
- Restricted-entry interval (REI) is 12 hours.

**Produced For:**

**ARYSTA LIFESCIENCE NORTH AMERICA, LLC  
15401 Weston Parkway, Suite 150  
Cary, NC 27513**

Arysta LifeScience and the Arysta LifeScience logo are registered trademarks of Arysta LifeScience Corporation. KANEMITE is a registered trademark of Agro-Kanesho Company.

The KANEMITE logo is a registered trademark of an Arysta LifeScience Group Company.

©2019 Arysta LifeScience Group Company.