

SUPPLEMENTAL LABELING

ACEQUINOCYL

GROUP

20B

ACARICIDE

KANEMITE® 15 SC MITICIDE

Supplemental Label to Add **DIRECTIONS FOR USE** on
BUSHBERRY (CROP SUBGROUP 13-07B)

THIS LABELING MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF APPLICATION. READ THE LABEL AFFIXED TO THE CONTAINER FOR **KANEMITE® 15 SC MITICIDE** BEFORE APPLYING. BOTH THE **KANEMITE 15 SC MITICIDE** CONTAINER LABEL AND THIS SUPPLEMENTAL LABELING MUST BE IN THE POSSESSION OF THE USER WHEN USING THE PRODUCT. USE OF **KANEMITE 15 SC MITICIDE** ACCORDING TO THIS LABELING IS SUBJECT TO THE USE PRECAUTIONS, RESTRICTIONS, AND LIMITATIONS IMPOSED BY THE LABEL AFFIXED TO THE CONTAINER FOR **KANEMITE 15 SC (EPA Reg. No. 66330-38)**.

This supplemental label expires on April 20, 2023 and must not be used or distributed after this date.

ACTIVE INGREDIENT:	% BY WT.
*Acequinocyl	15.8%
OTHER INGREDIENTS:	84.2%
TOTAL:	100.0%
*3-dodecyl-1, 4-dihydro-1, 4-dioxo-2-naphthyl-acetate.	
Contains 1.25 lb ai/gal.	

KEEP OUT OF REACH OF CHILDREN

CAUTION / PRECAUCIÓN

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

051520V001Su

BUSHBERRY (CROP SUBGROUP 13-07B): ARONIA BERRY; BLUEBERRY, HIGHBUSH; BLUEBERRY, LOWBUSH; BUFFALO CURRANT; CHILEAN GUAVA; CRANBERRY, HIGHBUSH; CURRANT, BLACK; CURRANT, RED; ELDERBERRY; EUROPEAN BARBERRY; GOOSEBERRY; HONEYSUCKLE, EDIBLE; HUCKLEBERRY; JOSTABERRY; JUNE BERRY (SASKATOON BERRY); LINGONBERRY; NATIVE CURRANT; SALAL; SEA BUCKTHORN; CULTIVARS, VARIETIES, AND/OR HYBRIDS OF THESE.

PEST	USE RATE
Broad mite (<i>Polyphagotarsonamus latus</i>)	31 fl oz/A
Southern red mite* (<i>Oligonychus ilicis</i>)	(0.3 lb ai/A)
Twospotted spider mite (<i>Tetranychus urticae</i>)	

RESTRICTIONS

- Do not apply this product through any type of irrigation system.
- Do not apply by air.
- Do not apply within 75 feet of aquatic areas.
- Not for indoor residential use.
- A 30-day plant-back interval is required for all crops not listed on this label.
- Apply in 50 to 100 gallons of water volume per acre.
- Allow a minimum of 21 days between applications.
- Do not make more than two applications of **KANEMITE 15 SC MITICIDE** per year.
- Do not apply more than 62 fl oz of **KANEMITE 15 SC MITICIDE** per acre (0.6 lb ai) per year.
- Preharvest interval is 1 day.
- Restricted-entry interval (REI) is 12 hours.
- Do not combine Kanemite 15 SC Miticide with Phos-ethyl (Fosetyl).

*Not registered for use in California.

Produced For:

ARYSTA LIFESCIENCE NORTH AMERICA, LLC

C/O UPL NA Inc.

630 Freedom Business Center, Suite 402

King of Prussia, PA 19406

Arysta LifeScience and the Arysta LifeScience logo are registered trademarks of a UPL Corporation Limited Group Company.

KANEMITE is a registered trademark of Agro-Kanesho Company.

The KANEMITE logo is a registered trademark of a UPL Corporation Limited Group Company.

©2020 UPL Corporation Limited Group Company