

FIFRA Section 24(c) Special Local Need Label



Dow AgroSciences

Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Kerb[®] SC

EPA Reg. No. 62719-578

SLN OR-110015

Weed Control in Sugarbeets Grown for Seed Only

For Distribution and Use Only in the State of Oregon

Expiration Date: This label is valid until December 31, 2022, or until otherwise amended, withdrawn, canceled, or suspended.

Active Ingredient:

pronamide: 3,5-dichloro-N-

(1,1-dimethyl-2-propynyl) benzamide.....35.6%

Other Ingredients64.4%

Total100.0%

Contains 3.3 lbs of active ingredient per gallon.

Keep Out of Reach of Children

CAUTION

Causes Moderate Eye Irritation.

Avoid contact with eyes or clothing.

Environmental Hazards

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

Directions for Use

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This Special Local Need (SLN) label and the federally registered label for this product must be in the possession of the user at the time of pesticide application.
- Follow all applicable directions, restrictions, and precautions on this SLN label and the EPA registered main label for Kerb SC, EPA Reg. No. 62719-578.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

General Information

Kerb[®] SC herbicide is a selective herbicide recommended for fall and winter applications to established seedling sugarbeets grown for seed for both preemergence and postemergence control of certain winter annual weeds and grasses.

Application Rate

Kerb SC herbicide should be applied at 2.5 pints (1.0 lb. active) per acre for preemergence or postemergence control of annual bluegrass, annual ryegrass, volunteer grains, and chickweed.

Kerb SC herbicide should be applied postemergence to weeds and sugarbeets during the fall or early winter months after the sugarbeets have three to five sets of true leaves. Kerb SC herbicide may be broadcast over the top of sugarbeets.

Mixing and Application

Mix the recommended amount of Kerb SC herbicide in water and apply in a minimum spray volume of 20 gallons per acre. Do not mix Kerb SC herbicide with spray adjuvants or pesticides other than Herbicide 273. Sprays should be applied to provide uniform distribution without skips or excessive overlapping of the spray pattern.

Kerb SC herbicide acts mainly through root absorption. In order to achieve good weed control, sufficient soil moisture from rain or irrigation is required following treatment to move Kerb SC herbicide into the soil to the weed root zone. Care should be taken to avoid spraying areas that may drain onto areas planted in cool season grasses or small grains.

Specific Use Restrictions

- **Chemigation:** Do not apply this product through any type of irrigation system.
- For use only on crops grown for seed as listed on this label in the state of Oregon.

Special Crop Use Restrictions

The pesticide applicator, the producer of the crop, and the seed conditioner must be aware that use of this product according to this labeling is deemed a non-feed/non-food use by the Oregon Department of Agriculture, and is regulated by Oregon Administrative Rule (OAR) 603-057-0535, Pesticide Use On Crops Grown for Seed. If the applicator of this pesticide is not the producer, the applicator must provide a copy of this labeling to the producer of the crop. Producers of this crop who use this product, or cause the product to be used on a field they operate, must provide a copy of this pesticide label to the seed conditioner.

This pesticide does not have an established pesticide residue tolerance for this crop. Consequently, no portion of this seed crop may be used or distributed for food or feed for one year (365 days) after the last application of this product. This restriction pertains to, but is not limited to, green chop, forage, hay, pellets, meal, whole seed, cracked seed, straw, roots, bulbs, foliage or seed screenings, and to the grazing of the crop field, stubble, or regrowth. All seed screenings shall be disposed of in such a manner that the screenings cannot be distributed or used for food or feed purposes, as indicated in OAR 603-057-0535. Additional regulations concerning seed screenings are stated in OAR 603-057-0535.

Any seed from a field treated with this pesticide product shall bear specific and conspicuous labeling on the container, or if shipped in bulk, on the shipment invoice or bill of lading. The labeling shall contain the following statement:

"This seed was produced using one or more products for which the United States Environmental Protection Agency has not established pesticide residue tolerances. This seed, in whole, as sprouts, or in any form, may not be used for human consumption or animal feed. Failure to comply with this condition may violate requirements of the Federal Food and Drug Administration, the Oregon Department of Agriculture and other regulatory agencies."

24(c) Registrant: Dow AgroSciences LLC
9330 Zionsville Road
Indianapolis, IN 46268-1054 USA
800-992-5994

®Trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow

R361-029
Accepted: 12/19/17
Replaces: R361-018