FIFRA Section 24(c) Special Local Need



PYRIPROXYFEN

GROUP

7C

INSECTICIDE

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF GEORGIA



INCREASE THE FREQUENCY OF APPLICATION WITH LOWER RATES FOR THE MANAGEMENT OF LEPIDOPTERA AND WHITEFLY IN BRASSICA HEAD AND STEM VEGETABLE GROUP, CUCURBIT VEGETABLES AND FRUITING VEGETABLES

This label for Knack[®] Insect Growth Regulator expires and must not be distributed or used in accordance with this SLN registration after December 31, 2027.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Follow all applicable directions, restrictions, worker protection standard requirements and precautions on the EPA registered label. This labeling must be in the possession of the user at the time of application.

CROPS AND COMMODITIES

Brassica Head and Stem Vegetable Group (Crop Group 5-16)

Broccoli; Brussels Sprouts; Cabbage; Cabbage, Chinese (napa); Cauliflower; Cultivars, varieties, and hybrids of these

Cucurbits (Crop Group 9)

Balsam Apple; Balsam Pear; Bitter Melon; Cantaloupe; Chayote; Chinese Cucumber; Chinese Waxgourd; Citron Melon; Cucumber; Edible Gourd; Gherkin; Momordica spp.; Muskmelon; Pumpkin; Summer Squash; Watermelon; Winter Squash

Fruiting Vegetables

African Eggplant; Bush Tomato; Bell Pepper; Cocona; Currant Tomato; Eggplant; Garden Huckleberry; Goji Berry; Ground Cherry; Martynia; Naranjilla; Okra; Pea Eggplant; Pepino; Pepper, Bell; Pepper, Nonbell; Roselle; Scarlet Eggplant; Sunberry; Tomatillo; Tomato; Tree Tomato; Cultivars, varieties, and/or hybrids of these

Pests	Application Rate	Special Instructions
Lepidoptera Beet Armyworm (suppression) Cabbage Looper (suppression) Diamond Back Moth (suppression) Imported Cabbage Worm (suppression) Western Yellow Stripped	5 fl oz/A (0.034 lb ai/A)	Apply Knack® Insect Growth Regulator when target pest(s) are beginning to establish or colonize in the field and/or before recommended threshold populations are observed. Do not wait until worms and/or caterpillar populations have exceeded their established thresholds. Apply by ground and in sufficient volume to obtain thorough coverage. Spray volume will depend on crop's canopy density (Use 20 to 50 GPA).
Armyworm (suppression)		

continued

GA-170005.R1 Page 1 of 2 R&RA 04/15/2022

Pests	Application Rate	Special Instructions
Whitefly Banded Wing Whitefly Greenhouse Whitefly Silverleaf Whitefly Sweetpotato Whitefly	5 fl oz/A (0.034 lb ai/A)	Knack Insect Growth Regulator must be part of a whitefly insecticidal program that utilizes different classes of chemistry implementing a rotation approach. Knack Insect Growth Regulator applications must be alternated with other insecticides and can only be applied sequentially if a whitefly insecticide is a tank mix partner.
		Timing : Apply first application when whitefly populations are starting to build, then make subsequent applications 7-14 days apart depending on whitefly migrations into the field and colonization.
		Knack Insect Growth Regulator must be rotated with other insecticide treatments targeting whiteflies with different modes of actions to reduce the risk of generating resistance among whitefly populations.
		Knack Insect Growth Regulator is not an adulticide and therefore an adulticide product is needed for control of Whitefly adults.

Use Restrictions

- Make applications every 7 to 14 days after last *Knack* Insect Growth Regulator treatment.
- Do not exceed 20 fl oz of *Knack* Insect Growth Regulator per acre per year.
- Regardless of formulation do not exceed 0.134 lb pyriproxyfen per acre per season.

Please contact Valent U.S.A. LLC at 800-6-VALENT (682-5368) if you have questions regarding use of this product.

FIFRA §24(c) Registrant: Valent U.S.A. LLC

P.O. Box 5075

San Ramon CA 94583

Knack is a registered trademark of Valent U.S.A. LLC

GA-170005.R1 Page 2 of 2 R&RA 04/15/2022