

FIFRA Section 24(c) Special Local Need



FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF GEORGIA



EPA Reg. No. 59639-95

SPLIT APPLICATION FOR CONTROL OF WHITEFLY IN VEGETATIVE COTTON
EPA SLN No. GA-210002

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Product	Rate & Timing
Knack® Insect Growth Regulator	5.0 fl oz/A (followed by) 5.0 fl oz/A
	Timing: Apply first application when whitefly populations are starting to build, then make second application 14 days later.
Vegetative Cotton: For vegetative (growing) cotton, it is critical the first application be made in a timely manner, prior to exponential population growth. A timely first application will keep populations low while a second application will maintain low populations while protecting new growth. Initiate first spray when 5 th leaf below the terminal averages 3-5 adults and at least one large nymph per sample. If adult populations are higher, add an adulticide for knockdown.	
Non-Vegetative Cotton: If cotton growth is no longer vegetative or “cut out”, make one application of <i>Knack</i> at 8 to 10 fl oz/A, again when populations are low. Add an adulticide for knockdown.	

Use Restrictions

- Always read and follow all label directions when using any pesticide alone or in tank mix combinations. The most restrictive labeling applies when using a tank mix.
- Do not apply more than 10 oz/A of *Knack* per season.
- Do not apply within 28 days of harvest.
- Refer to labels for additional restrictions and use directions.

This label and the EPA registered label must be in the possession of the user at the time of mixing and pesticide application.

Follow all applicable directions, restrictions and precautions on the EPA-registered labels.

THIS RECOMMENDATION EXPIRES DECEMBER 31, 2025

FIFRA Section 24(c) Registrant: Valent USA LLC
P.O. Box 5075
San Ramon, CA 94583

Knack is a registered trademark of Valent U.S.A. LLC