Recommendation

RESTRICTED USE PESTICIDE
DUE TO TOXICITY TO FISH AND AQUATIC ORGANISMS
For retail sale to and use only by Certified Applicators, or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.

FIFRA SECTION 2(ee) RECOMMENDATION FOR DISTRIBUTION AND USE IN ALABAMA, DELAWARE, FLORIDA, GEORGIA, LOUISIANA, MARYLAND, MISSISSIPPI, NEW JERSEY, NORTH CAROLINA, PENNSYLVANIA, SOUTH CAROLINA, TENNESSEE AND VIRGINIA

Lambda-Cy™ EC Insecticide-RUP
EPA Reg. No. 70506-121

For control of Kudzu bug

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by US EPA.

DIRECTIONS FOR USE
It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read and follow all applicable instructions, restrictions, and precautions on the EPA-approved label. This label and the federally registered label must be in the possession of the user at the time of pesticide application.

CROP	RATE LAMBDA-CY EC INSECTICIDE per ACRE	REMARKS
Legume Vegetables: Soybeans	3.84 fl. oz. (0.03 lbs ai/A)	Do not apply within 30 days of harvest. Do not apply more than 0.06 lb. ai (0.48 pt)/A per season. Do not graze or harvest treated soybean forage, straw or hay for livestock feed. Apply by ground or air using enough water to obtain full coverage of foliage. When applying by air, apply in at least 2 gals. of water/A. Use scouting to determine need for application, usually at intervals of 5 or more days. Base the timing and frequency of applications on when insect populations reach local economic thresholds.

United Phosphorus, Inc.
630 Freedom Business Center, Suite 402
King of Prussia, PA 19406
1-800-438-6071

Recommendation

CROP	RATE LAMBDA-CY EC INSECTICIDE per ACRE	REMARKS
Legume Vegetables: Beans and Peas	3.84 fl. oz. (0.03 lbs ai/A)	For edible podded and succulent shelled legume vegetables, do not apply within 7 days of harvest. For dried shelled legume vegetables, do not apply within 21 days of harvest. Do not apply more than 0.12 lb. ai (0.96 pt. or 15.36 fl. oz. of product)/A per season. For succulent and dried shelled peas and beans, do not graze livestock in treated areas or harvest vines for forage or hay. Apply by ground or air using enough water to obtain full coverage of foliage. When applying by air, apply in at least 2 gals. of water/A. Use scouting to determine need for application, usually at intervals of 5 or more days. Base the timing and frequency of applications on when insect populations reach local economic thresholds.

Rev. 08/05/2013

United Phosphorus, Inc.
630 Freedom Business Center, Suite 402
King of Prussia, PA 19406
1-800-438-6071