



β -CYFLUTHRIN	GROUP	3	INSECTICIDE
IMIDACLOPRID	GROUP	4A	INSECTICIDE

RESTRICTED USE PESTICIDE

Due to Toxicity to Fish and Aquatic Organisms

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

FIFRA 2(ee) Recommendation

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE(S) OF
ARIZONA AND CALIFORNIA

LEVERAGE[®] 360 Insecticide

EPA Reg. No. 264-1104

For control of brown marmorated stink bug in Citrus, Grape and Potato

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. This product bulletin should be in the possession of the user at the time of application of the product. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its label.

CITRUS (California and Arizona, Only)

Crops of Crop Group 10 Including: Calamondin, Citrus citron, Citrus hybrids (*Citrus* spp., includes chironja, tangelo and tangor), Grapefruit, Kumquat, Lemon, Lime, Mandarin (tangerine), Orange (sweet and sour), Pummelo, Satsuma mandarin, and other cultivars and/or hybrids of these

Pests Controlled	Product Rate fluid ounces/Acre
Brown Marmorated Stink Bug	3.2 – 6.4

RESTRICTIONS

Pre-Harvest Interval (PHI): **0 day**

Minimum interval between applications: **10 days**

Maximum LEVERAGE 360 Insecticide allowed per year: **6.4 fluid ounces/Acre** (0.05 lb AI/A β -cyfluthrin, 0.10 lb AI/A imidacloprid)

Maximum β -cyfluthrin allowed in all forms per crop season: 0.05 lb AI/A

Maximum cyfluthrin allowed in all forms per crop season: 0.10 lb AI/A

Maximum β -cyfluthrin plus cyfluthrin allowed in all forms per crop season: 0.10 lb AI/A

Do not apply during bloom or within 10 days prior to bloom or when bees are foraging.

LEVERAGE 360 Insecticide is to be applied through properly calibrated ground or aerial application equipment.

Minimum application volume (water): 25 GPA – ground, 25 GPA – aerial application.

ADDITIONAL USE DIRECTIONS

Time applications for control of scale according to crawler stage. Two applications may be required to achieve control.

β-CYFLUTHRIN	GROUP	3	INSECTICIDE
IMIDACLOPRID	GROUP	4A	INSECTICIDE

GRAPE Includes: Table grape, Raisin, Wine and Muscadine grape	
Pests Controlled	Product Rate fluid ounces/Acre
Brown Marmorated Stink Bug	2.8 – 3.2
RESTRICTIONS Pre-Harvest Interval (PHI): 3 days Minimum interval between applications: 14 days Maximum LEVERAGE 360 Insecticide allowed per year: 6.4 fluid ounces/Acre (0.05 lb AI/A β-cyfluthrin, 0.10 lb AI/A imidacloprid) Maximum β-cyfluthrin allowed in all forms per crop season: 0.1 lb AI/A Maximum cyfluthrin allowed in all forms per crop season: 0.2 lb AI/A Maximum β-cyfluthrin plus cyfluthrin allowed in all forms per crop season: 0.2 lb AI/A Minimum application volume (water): 25 GPA – ground, 10 GPA – aerial application. ADDITIONAL USE DIRECTIONS LEVERAGE 360 Insecticide is to be applied through properly calibrated application equipment.	

POTATO	
Pests Controlled	Product Rate fluid ounces/Acre
Brown Marmorated Stink Bug	2.8
RESTRICTIONS Pre-Harvest Interval (PHI): 7 days Minimum interval between applications: 7 days Maximum LEVERAGE 360 Insecticide allowed per year: 12.8 fluid ounces/A (0.1 lb AI/A β-cyfluthrin, 0.2 lb AI/A imidacloprid) Maximum β-cyfluthrin allowed in all forms per crop season: 0.132 lb AI/A Maximum cyfluthrin allowed in all forms per crop season: 0.263 lb AI/A Maximum β-cyfluthrin plus cyfluthrin allowed in all forms per crop season: 0.263 lb AI/A ADDITIONAL USE DIRECTIONS LEVERAGE 360 Insecticide is to be applied through properly calibrated ground, aerial or chemigation application equipment. ^{1/} For all aphids, apply as pest population begins to build and prior to buildup of damaging levels. For aphid control in crop with dense canopy use ground application equipment which will provide thorough coverage of lower leaves.	

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577.

For **PRODUCT USE** Information Call 1-866-99BAYER (1-866-992-2937).

As with any crop-protection product, always read and follow label directions.

For additional information call toll-free 1-866-99BAYER (1-866-992-2937).

Bayer CropScience LP
800 N. Lindbergh Blvd.
St. Louis, MO 63167