



# Liquid Zinc 9%

## MICRONUTRIENT COMPLEX

Read Entire Label Before Using This Product

### GUARANTEED ANALYSIS

Sulfur (S).....	4.5%
Zinc (Zn).....	9.0%

Derived from zinc sulfate and lignin sulfonate.

### GENERAL INFORMATION

LIQUID ZINC 9% is especially recommended for citrus crops but may also be used on field, row, vegetable, fruit, tree, vine, ornamental and turf crops. LIQUID ZINC 9% alone, with pesticides, or with other fertilizers, may be applied directly to the soil or to the foliage of growing crops. LIQUID ZINC 9% may be applied via ground sprayer (high volume or low-volume), aircraft or drip irrigation systems.

### DIRECTIONS FOR USE

Soil applications of LIQUID ZINC 9% are recommended with nitrogen fertilizer solutions. Thorough coverage and wetting are needed for optimum foliar spray results. Actively growing plants with young leaves absorb foliar applied LIQUID ZINC 9% most efficiently. Immediate postirrigation or post-rainfall application gives the best foliar uptake. Addition of urea (for example, 5 lbs. Lo-Bi Urea per 100 gallons of spray solution) will enhance foliar uptake of LIQUID ZINC 9%.

**COMPATIBILITY:** LIQUID ZINC 9% is compatible with most pesticides and fertilizers. However, if the desired combination has not been previously used, a compatibility test is recommended.

**MIXING:** Shake, stir or swirl contents before using. Always add this product to the spray tank before adding pesticides.

### FOLIAR APPLICATION - ALL CROPS:

LIQUID ZINC 9% applications should be a minimum of 7-14 days apart. Apply 2-4 quarts/acre as needed on a 7-14 day schedule. AVOID foliar applications of LIQUID ZINC 9% when plants are under moisture stress. AVOID foliar applications of LIQUID ZINC 9% to deciduous tree crops during bloom.

**Dilute Applications:** Apply the suggested dose of product in enough water for thorough coverage.

**Concentrate Applications:** Use up to 2 quarts of LIQUID ZINC 9% in a minimum of 25 gallons of water.

**Aerial Applications:** Up to 1 quart of LIQUID ZINC 9% per gallon of spray may be used, but more dilute solutions will improve coverage and results.

### SOIL APPLICATION - ALL CROPS\*

Soil Fertility Maintenance .....	1-3 qts./acre
Mild Deficiency .....	2-4 qts./acre
Severe Deficiency.....	1-3 gal./acre

\* The lower rates are for field, row and vegetable crops and the higher rates are for fruit, tree, vine, ornamental and turf crops.

**Fertigation:** Apply the recommended amount of LIQUID ZINC 9% in split applications one to two weeks apart throughout the growing season.

Information regarding the contents and levels of metals in this product is available by calling 1-800-696-6108.



DANGER: Causes serious eye damage. May cause cancer.

**Precautionary Statements:** Obtain special instructions before use. Do not handle until all safety precautions have been read and understood. Wear eye/face protection and protective gloves/clothing. Avoid release to the environment. Collect spillage. **First Aid:** *If in eyes:* Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. Immediately call a doctor. **Storage and Disposal:** Do not contaminate water, food or feed by storage or disposal. Keep out of reach of children. Store locked up. Dispose of contents/container in accordance with local regulations.

**CONDITIONS OF SALE AND WARRANTY:** GAR Bennett LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated when used in accordance with the use directions under normal conditions. GAR Bennett LLC neither makes, nor authorizes any agent or representative to make, any other warranties, express or implied, including fitness or merchantability. The directions for use of this product are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as timing and method of application, weather and crop conditions, presence of other materials, or other influencing factors, all of which are beyond the control of GAR Bennett LLC. Buyer and user acknowledge and assume all risks and liability resulting from the handling, storage and use of this material not in strict accordance with directions given herewith. In no case shall GAR Bennett LLC be liable for consequential, special, indirect, or incidental damages or losses resulting from the handling or use of this product. The foregoing is a condition of sale by GAR Bennett LLC and is accepted as such by the Buyer.

**WARNING:** Cancer and Reproductive Harm - [www.P65Warnings.ca.gov](http://www.P65Warnings.ca.gov)

Manufactured for:

Proud Owner Of:



Integrated  
Agribusiness  
Professionals

Gar Bennett LLC

8246 S. Crawford Avenue - Reedley, CA 93654  
(559) 638-6311 tel (559) 638-6314 fax  
[www.garbennett.com](http://www.garbennett.com)

Net Weight 11.0 lbs/gallon @ 68° F

Net Contents: Bulk \_\_\_\_\_ gallons

01/19