GENERAL INFORMATION

LOAD contains available forms of the essential nutrients nitrogen, boron, and molybdenum, and is non-phytotoxic when used as directed.

DIRECTIONS FOR USE

It may be a violation of law to use this product inconsistent with its labeling.

LOAD can be applied to the plant foliage by spraying. Apply the recommended amount in a volume sufficient to provide uniform coveage.

RECOMMENDED USES:

FOR FOLIAR APPLICATION: Apply LOAD to the foliage, every 7 to 14 days, at 0.5 to 2 quarts per acre on field crops, 1 to 4 quarts per acre on vegetable, fruit, and nut crops.

Use **LOAD** at the recommended rate on the following crops.

FIELD CROPS: Alfalfa, bean crops, canola, clover, corn, cotton, flax, forage grasses, peanut, potato, rape (canola), rice, sorghum, soybean, sugar beet, sugarcane, sunflower, tobacco, and wheat, barley, oat, or other cereal grain. Apply prior to flowering, early flowering, and within 7-14 days' post flowering/early seed development. On alfalfa apply within 7 days after cutting.

VEGETABLE CROPS: Artichoke, asparagus, broccoli, cauliflower, cabbage, kale, and other cruciferous vegetables; leaf lettuce, head lettuce, dandelion, radicchio, and other leafy vegetables; cucumber, melon, pumpkin, squash, and other cucurbits, onion, garlic, and other bulb vegetables; tomato, pepper, eggplant and other fruiting vegetables; strawberry; carrot, sweet potato, radish, beet and other root and tuber vegetables. Apply just prior to or at early flowering.

FRUIT AND NUT CROPS: Apple, avocado, blueberry and other bush berries, blackberry, raspberry and other cane berries, sweet orange, lemon, mandarin and other citrus, date palm, fig, grapes, hops, oil palm, olive, persimmon, cherry, peach, plums, prune, nectarine, apricot and other stone fruit, almond, pecan, pistachio, walnut and other tree nuts, pear, pomegranate, and subtropical fruit. Apply at early flowering, at petal fall, and early fruit development.

MIXING INSTRUCTION: Follow this mixing order 1. Water 2. **LOAD** 3. Pesticide. **LOAD** will disperse in water with little agitation. LOAD is compatible with most fertilizers and pesticides. Always conduct a jar test when using new or untried combinations.

Notice: The information contained herein is true and accurate to the best of our knowledge and belief, but is presented without guarantee since field conditions and use are beyond our control. Neither the manufacturer nor the seller make any warranty, expressed or implied concerning the use of this product other than indicated on the label. Buyer assumes all risk of use of this material when such use is contrary to label instruction. Read and follow the label directions carefully.

CONDITIONS OF SALE: 1. Seller warrants that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use. No one, other than an officer of Seller, is authorized to make any warranty, guarantee of direction concerning this product. 2. Because of time, place, rate of application and other conditions of use are beyond sellers control, Sellers liability from handling, storage and use of this product is limited to replacement of product or refund of purchase price.



2-0-0

Derived from boron ethanolamine and sodium molybdate

CAUTION

KEEP OUT OF REACH OF CHILDREN; MAY BE HARMFUL IF SWALLOWED, INHALED OR IN CONTACT WITH SKIN; MAY CAUSE EYE IRRITATION. MAY CAUSE DAMAGE THROUGH PROLONGED EXPOSURE.

PRECAUTIONS: Do not breathe dust/fume/gas/mist/vapors/spray. Wash hands thoroughly after handling. Do not eat, drink or smoke when using this product. Use only outdoors or in well-ventilated area. Avoid release to environment. Wear protective gloves/protective clothing/eye protection/face protection. Keep container tightly closed. Keep cool and protected from sunlight. If medical advice is needed, have product container or label at hand. For more details regarding the safe use/handling, storage and disposal of this product as well as additional hazards and first aid measures, please obtain the current Safety Data Sheet at www.stollerusa.com.

STORAGE AND DISPOSAL

We recommend storing pails a maximum of three pails high. Do not stack pallets. Store container tightly closed in a cool/well-ventilated dry place protected from sunlight above 40°F. Do not allow container to freeze. Store locked up. DISPOSE of contents/ container at a permitted facility or as advised by your local regulatory authority. Do not contaminate water, food or feed by storage or disposal. Do not apply directly to areas where surface water is present, or to intertidal areas below the mean high water mark.Do not contaminate water when disposing of equipment wash water. Do not reuse empty container.

WARNING: This product contains boron at a level that may be detrimental to growing plants. Use of boron on crops other than those recommended may result in serious injury to those crops.

WARNING: The application of fertilizing materials containing molybdenum may result in forage crops containing levels of molybdenum which are toxic to ruminant animals.

Information regarding the contents and levels of metals in this product is available on the internet at http://www.aapfco.org/metals.html



2-0-0

2399

Density: 10.2 lb/gal (1.22 kg/L) @ 68°F (20°C)

NET CONTENT	NET WEIGHT
☐ 2.50 gal (9.5 L)	25.5 lb (11.5 kg)
☐ 2.64 gal (10.0 L)	26.9 lb (12.2 kg)
☐ 5.00 gal (18.9 L)	51.0 lb (23.1 kg)
☐ 5.28 gal (20.0 L)	53.9 lb (24.4 kg)
☐ 55.00 gal (208.2 L)	561.0 lb (254.0 kg)
☐ 275.00 gal (1041.0 L)	2805.0 lb (1270.0 kg)

☐ BULK



Manufactured and Guaranteed by Stoller Enterprises, Inc.

9090 Katy Freeway, Suite 400 Houston, TX 77024 Phone: 1-800-539-5283 or 713-461-1493



