

# FIFRA 24(c)

## Special Local Need

Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

### RESTRICTED USE PESTICIDE

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

## Lorsban<sup>®</sup> Advanced

EPA Reg. No. 62719-591

EPA 24(c) Special Local Need Registration SLN LA-150004  
For Distribution and Use Only in the State of Louisiana

**Insect Control Infesting Sweet Potato (60-Day PHI)**

### ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Lorsban<sup>®</sup> Advanced insecticide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Lorsban Advanced according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Lorsban Advanced.

### Directions for Use

Refer to product label for Lorsban Advanced for Use Precautions and Restrictions, Mixing and Application instructions.

Target Pests	Lorsban Advanced (pint/acre)
Conderus (wireworm) sweet potato flea beetle Systema (flea beetle)	4

Apply to the soil surface as a preplant broadcast spray to reduce the feeding damage caused by listed pests. Use a spray volume of 10 gallons or more per acre. Incorporate immediately after application to a depth of 4 to 6 inches using a rotary hoe, disc cultivator, or other suitable incorporation equipment. Plant sweet potatoes in the usual manner no more than 14 days after treatment. Delaying planting more than 14 days after application will reduce the time interval of protection against feeding damage.

### Specific Use Precautions and Restrictions:

- **Worker Restricted Entry Interval:** Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours unless PPE required for early entry is worn.
- **Preharvest Interval:** Do not harvest within 60 days of treatment.
- Lorsban Advanced will not control false wireworms, white fringe beetle or other grubs that attack sweet potatoes.
- Do not make more than one application of Lorsban Advanced or other product containing chlorpyrifos per season.
- Maximum single application rate is 1.88 lb ai chlorpyrifos per acre.
- **Expiration Date: March 20, 2025**

<sup>®</sup>™ Trademarks of Dow AgroSciences, DuPont or Pioneer and their affiliated companies or respective owners

R368-136

Approved: 03/20/2020

Replaces R368-103