



Bayer CropScience LP  
 P.O. Box 12014  
 2 T.W. Alexander Drive  
 Research Triangle Park, North Carolina 27709  
 1-866-99BAYER (1-866-992-2937)

## Luna<sup>®</sup> Experience

For control of shot hole, jacket rot, green fruit rot, and peach leaf curl in cherries, peaches, and plums

EPA Reg. No. 264-1091

## FIFRA 2(ee) Recommendation

### FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE(S) OF CALIFORNIA

**FIFRA Section 2(ee) Recommendation:** This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. This product bulletin should be in the possession of the user at the time of application of the product. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

#### DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its label.

#### Cherry subgroup 12-12A; Peach subgroup 12-12B; and Plum subgroup 12-12C

**Cherry subgroup:** Capulin; Cherry, black; Cherry, Nanking; Cherry, sweet; Cherry, tart; cultivars, varieties, and/or hybrids of these.

**Peach subgroup:** Peach; Nectarine; cultivars, varieties, and/or hybrids of these.

**Plum subgroup:** Apricot; Apricot, Japanese; Jujube, Chinese; Plum; Plum, American; Plum, beach; Plum, Canada; Plum, cherry; Plum, Chickasaw; Plum, Damson; Plum, Japanese; Plum, Klamath; Plumcot; Plum, prune; Sloe; cultivars, varieties, and/or hybrids of these.

Disease Control	Application Rate	Product Instructions
Shot hole ( <i>Wilsonomyces carpophilus</i> )	6.0 to 10 fl oz/acre	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 7- to 14-day interval. When disease pressure is severe, use the higher rates and/or shorter intervals.
Jacket rot, Green fruit rot ( <i>Botrytis cinerea</i> )		
Peach leaf curl ( <i>Taphrina</i> spp.)		

#### Restrictions:

- Do not apply more than 34 fl oz of LUNA EXPERIENCE per acre per year.
- Apply using ground, aerial, or chemigation equipment.
- Regardless of formulation or method of application, do not apply more than 0.446 lbs Fluopyram or 1.35 lbs tebuconazole per acre per year, including soil and foliar uses.
- May be applied the day of harvest. To limit the potential for development of disease resistance to these fungicide classes, do not make more than 2 sequential applications of LUNA EXPERIENCE or any Group 7 or Group 3 containing fungicide before rotating with a fungicide from a different Group.

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577.  
For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937).

*As with any crop-protection product, always read and follow label directions.  
For additional information call toll-free 1-866-99BAYER (1-866-992-2937).*