



Bayer CropScience LP  
P.O. Box 12014  
2 T.W. Alexander Drive  
Research Triangle Park, North Carolina 27709  
1-866-99BAYER (1-866-992-2937)

## LUNA<sup>®</sup> EXPERIENCE

Reduced Application Rate for use in  
watermelon

EPA Reg. No. 264-1091

### PRODUCT BULLETIN

**FOR DISTRIBUTION AND USE ONLY WITHIN THE STATES OF ARIZONA, ARKANSAS, CALIFORNIA, COLORADO, CONNECTICUT, DELAWARE, FLORIDA, GEORGIA, IDAHO, ILLINOIS, INDIANA, IOWA, KENTUCKY, MAINE, MARYLAND, MICHIGAN, MINNESOTA, MISSISSIPPI, MISSOURI, MONTANA, NEBRASKA, NEW JERSEY, NEW MEXICO, NORTH CAROLINA, NORTH DAKOTA, OHIO, OKLAHOMA, OREGON, PENNSYLVANIA, RHODE ISLAND, SOUTH CAROLINA, TENNESSEE, TEXAS, UTAH, VERMONT, VIRGINIA, WASHINGTON, WEST VIRGINIA, WISCONSIN**

**FIFRA Section 2(ee) Recommendation:** This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. It is a violation of federal law to use this product in a manner inconsistent with its label. This product bulletin should be in the possession of the user at the time of application of the product. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

#### DIRECTIONS FOR USE

**WATERMELON:** (includes hybrids and/or varieties of *Citrullus lanatus*).

Disease Control	Product Rate	Use Information
Gummy stem blight ( <i>Didymella bryoniae</i> )  Belly rot ( <i>Rhizoctonia solani</i> )  Anthracnose ( <i>Colletotrichum spp.</i> )	10.0 to 17 fl oz/acre	Begin fungicide applications preventatively. Continue as needed on a 10- to 14-day interval. When disease pressure is severe, use the higher rates and/or shorter intervals.
Powdery mildew ( <i>Sphaerotheca fuliginea</i> / <i>Podosphaera xanthii</i> ) ( <i>Erysiphe cichoracearum</i> )	6.0 to 17 fl oz/acre	Begin fungicide applications preventatively. Continue as needed on a 10- to 14-day interval. When disease pressure is severe, use the higher rates and/or shorter intervals.
Alternaria leaf spot ( <i>Alternaria cucumerina</i> )	8 to 17 fl oz/acre	Begin fungicide applications preventatively. Continue as needed on a 10- to 14-day interval. When disease pressure is severe, use the higher rates and/or shorter intervals.

**Note:** A mild yellowing on leaf margins is sometimes seen following application of LUNA<sup>®</sup> EXPERIENCE in watermelon.

**Restrictions:** Do not apply more than 34 fl oz of LUNA<sup>®</sup> EXPERIENCE per acre per season. Do not apply LUNA<sup>®</sup> EXPERIENCE within 7 days of harvest. May not be applied with aerial application equipment. To limit the potential for development of disease resistance to these fungicide classes, do not make more than 2 sequential applications of LUNA<sup>®</sup> EXPERIENCE or any Group 7 or Group 3 containing fungicide before rotating with a fungicide from a different Group.

Use of this product according to this bulletin has not been reviewed or endorsed by the State of Indiana.

For **MEDICAL** And **TRANSPORTATION** Emergencies **ONLY** Call 24 Hours A Day 1-800-334-7577  
For **PRODUCT USE** Information Call 1-866-99BAYER (1-866-992-2937)

As with any crop-protection product, always read and follow label directions.  
For additional information call toll-free 1-866-99BAYER (1-866-992-2937).