

The Go To Company

P.O. Box 5569 ▲ Yuma, AZ 85366-5569 ▲ Phone (928) 783-8844 ▲ FAX (928) 343-9255



EPA Reg. No. 10163-324

2(ee) Recommendation

For use in the State of Alabama, Arizona, Arkansas, California, Colorado, Connecticut, Delaware, District of Columbia, Florida, Georgia, Hawaii, Idaho, Illinois. Indiana, Iowa, Kansas, Kentucky, Louisiana, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Utah, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming

Insecticide/Miticide/Fungicide for Control of Whitefly & Mites
This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or
approved by EPA.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product inconsistent with its labeling.

All applicable directions, restrictions, and precautions on the EPA registered label are to be followed.

This labeling must be in the possession of the user at the time of pesticide application.

Do not apply this product in any way that will contact workers or other persons; either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe; consult the agency responsible for pesticide regulation. Always read this FIFRA 2(ee) bulletin and the entire label on the product container or booklet attached before using this product.

PRODUCT INFORMATION

M-Pede® insecticide/fungicide is a contact insecticide, miticide and fungicide for control of soft-bodied insects, mites and powdery mildew. The formulation is based on potassium salts of naturally derived fatty acids. This product may be used to control targeted pests on crops which include vegetables, grapes and other small fruits, tree fruits, tree nuts, cotton, tobacco, shrubs, shade and ornamental trees, turf, foliage and flowering plants growing outdoors, in greenhouses and in interiorscapes. This product can be applied up to harvest.

Pest	Requirements
Whitefly & mites* at egg and larval stages	Stand Alone Rate : as a pre-plant dip, apply 0.25-0.5% v/v solution of this product to ornamental transplants. Always determine the sensitivity before applying to new seedlings and transplants. Tank Mix Rate : as a pre-plant dip, apply 0.5% v/v solution of this product in combination with a recommended product to ornamental transplants. Always determine the sensitivity before applying to new seedlings and transplants.
*Not for use in the state of New York	

Restrictions:

- Do not make repeat applications at less than 7-day intervals, or make more than 3 sequential applications at 7- to 14-day intervals unless your experience has shown the use pattern to be non-injurious to the type and the development stage of plant(s) to be treated under local growing conditions
- Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.
- Always determine the compatibility of water, pesticides and other tank additives with this product.
- Do not dip stressed cuttings or transplants.