RECOMMENDATION FOR USE:			
Crop	Rate Pints /Acre	Rate Liters /HA	Number & Timing of Application
Almonds	3-4	3.5-5.0	Make 3 to 4 applications from bud swell through jacket split
Apples	3-6	3.5-5.0	Make 3 to 5 applications, start at post thinning through early sizing, including 14 to 21 days post harvest.
Apricots, Nectarines Peaches	3-4	3.5-5.0	Make 3 applications, first at bud swell, second after jacket split and third 30-45 days pre harvest.
Avocados	3-6	3.5-7.0	Multiple applications as required. Do not apply within 30 days of harvest.
Broccoli	3-4	3.5-5.0	4-6 applications starting shortly before head formation.
Cabbage Cauliflower Lettuce	3-4	3.5-5.0	4-6 applications starting shortly before head formation.
Cane Berries	3-4	3.5-5.0	Make 2 applications, 1st immediately post bloom, 2nd 21 to 30 days later.
Celery Chicory	3-4	3.5-5.0	Weekly applications starting before black heart symptoms arise.
Cherries, Plums, Prunes	3-4	3.5-5.0	Make 3 to 4 applications, 1st at bud swell, 2nd at petal fall, 3rd at first color, 4th 30 to 45 days pre-harvest.
Citrus	3-6	3.5-7.0	Multiple applications as required.
Cotton	5	6.0	3 applications between 5-7 leaf stage.
Cucumber, Melon, Pepper, Tomato	1-3	1.5-3.5	Make 6 to 12 applications as needed during fruit development.
Cut Flowers	1-2		Apply 1-2 pints per 1,000 sq. ft. every 10 to 14 days as needed.
Grapes	3-6		Make 2 to 3 applications, start post shatter through early ripening.
Hazelnuts	3-8	3.5-9	Make 3 to 4 applications, start at bud swell through early development.
Kiwi	3-4	3.5-5.0	Make 2 to 3 applications, start at early fruit formation to ripening.
Omamental Woody Plants	2-4		Make multiple applications during growing season, begin at first growth through early fall.
Pears	3-6	3.5-7.0	Make 3 to 4 applications, 1st at bud swell, 2nd 1 to 2 weeks after petal fall, 3rd at early fruit sizing, 4th optional 40-45 days pre-harvest.
Pistachios	3-6	3.5-7.0	Make 3 to 4 applications, 1st at bud swell, 2nd at green tip, 3rd at jacket split, 4th optional if required.
Potatoes	2-4	2.5-5.0	Multiple applications during period of heat stress.
Table Grape	3-4	3.5-5.0	Make 2-3 applications, start at bloom through ripening as required
Strawberries	4-5		Make 3 applications in conjunction with last 3 pre-harvest pesticide sprays
Walnuts	3-6		Make 2-3 applications, 1st at bud swell, 2nd at catkin elongation, 3rd at first set

GROW MORE

MAXI-GAL

Suspension

10% Calcium

with Magnesium &

7 Micronutrients

Export Made In U.S.A.





Ĉa

DIRECTIONS

Grow More MAXI-CAL is a water soluble foliar fertilizer designed to provide Nitrogen enhanced secondary and micronutrients for agricultural crops. May be applied by itself or in combination with pesticides.

Rates are based on spray volumes of 50-150 gallons/acre (500-1,500 liters/hectare) For higher volume sprays, rates should be adjusted.

Pre-dilute Grow More MAXI-CAL in water prior to adding to spray tank. Add MAXI-CAL to spray tank when half-full, while agitating, prior to adding pesticide. Spray immediately after mixing. Do not store solution mixed with pesticide.

CAUTION

Store MAXI-CAL in temperatures between 41°F to 104°F and avoid extreme variations in temperature. If separation occurs during long termed storage, shake vigorously before using.

CAUTION: KEEP OUT OF REACH OF CHILDREN

The seller hereby represents and warrants that the fertilizers shall be manufactured and shipped to purchaser in a form suitable for application to plants, trees and other agricultural crops and hereby further represents and warrants that the fertilizer products shall meet the guaranteed minimum analysis as set forth on the product's identity labels. However, this warranty does not extend to the use of these products contrary to label instructions and the buyer assumes the risk of any such use.

GUARANTEED ANALYSIS 10-0-0

Total Nitrogen10.00%
8.76% Nitrate Nitrogen
0.54% Urea Nitrogen
Calcium10.00%
Magnesium (Mg)1.10%
Boron (B)0.02%
Colbalt (Co)
Copper (Cu)0.05%
0.05% Chelated Copper
Iron (Fe)0.10%
0.10% Chelated Iron
Manganese (Mn)0.05%
0.05% Chelated Manganese
Molybdenum (Mo)0.0005%
Zinc (Zn)0.05%
0.05% Chelated Zinc

Derived from Calcium Nitrate, Calcium Ammoniuum Nitrate, Urea, Magnesium Nitrate, Boric Acid, Copper EDTA, Iron EDTA, Manganese EDTA, Zinc EDTA, Ammonium Molybdate and Cobalt Sulfate.

WARNING: Application of fertilizer materials containing Molybdenum may result in forage crops containing levels of Molybdenum which are toxic to ruminant animals. This product also contains Boron and may also result in injury to crops.

Information regarding the contents and levels of metals in this product is available by calling 1-800-338-7160

Manufactured By:

GROW MORE

15600 New Century Drive,
Gardena, CA, U.S.A. 90248

NET WT. 2.5 gallons 2X2.5 gallons Density 13lbs/gal.

Visit Us At: www.growmore.com