MAXI-PLEX

Multi-Purpose Micronutrient For Soil & Foliage Agriculture • Horticulture



5-0-0

GUARANTEED ANALYSIS

Total Nitrogen (N)	5.00%
Magnesium (Mg)	4.00%
1.00% Chelated Magnesium	
3.00% Water Soluble Magnesium	
Boron (B)	0.50%
Cobalt (Co)	0.05%
Copper (Cu)	1.50%
1.50% Chelated Copper	

Iron (Fe)	3.50%
3.50% Chelated Iron	
Manganese (Mn)	4.00%
4.00% Chelated Manganese	
Molybdenum (Mo)	0.10%
Zinc (Zn)	2.00%
2.00% Chelated Zinc	

Derived from Magnesium EDTA, Magnesium Sulfate, Boric Acid, Cobalt Sulfate, Copper EDTA, Iron EDTA, Manganese EDTA, Sodium Molybdate & Zinc EDTA.

GENERAL DIRECTIONS

Maxi-Plex may be applied to vegetable field and row crops, and fruit and nut trees, vines and citrus.

For best results apply what crops have adequate supplies of water and major nutrients.

DIRECTIONS FOR USE

Mix in water before use, add to tank when half full, agitate until dissolved. If pesticides are to be use add to tank last compatibility. If the intended combination of fertilizer or pesticide has not been previously use, a compatibility test is recommended.

Conditions that cause one particular nutrient deficiency can also induce deficiencies of other trace elements.

FOLIAR APPLICATION

Field vegetable, row crops use 0.50 to 1 lb. of MAXI-PLEX per acre (0.50 to 1 kg per HA) in sufficient water for full coverage, 1 lb. per 100 gallons of water.

Fruit and nut trees, vines, use 0.50 to 1 lb of MAXI-PLEX per acre (0.50 to 1 kg per HA) in sufficient water for full coverage spray, 1 lb. per 100 gallons of water.

AERIAL APPLICATION

Apply 1/3 lb. of MAXI-PLEX in at least 10 gallons of water. (150 grams in 40 liters)

CAUTION: Not recommended to apply more than 1 lb. of MAXI-PLEX per foliar application as crop injury may occur. Multiple application should be made in 10 to 14 day intervals to reduce the potential for crop injury.

SOIL APPLICATION

MAXI-PLEX may be used on all types of soil, dissolve in water or fluid fertilizer or water soluble fertilizer for band, sidedress or broadcast application. Soil application can also be made by dissolving in water then into fertigation system. Field, vegetable or row crops.

Use up to 12 lbs. per acre (13 kgs per HA)

Vines

0.50 to 1 lb. per 100 feet of row (225 to 500 grams per 30 meter row)

Fruit and nut trees:

1 to 2 lbs. per tree (400 to 900 grams per tree)

CAUTION: KEEP OUT OF REACH OF CHILDREN

The seller hereby represents and warrants that the fertilizers shall be manufactured and shipped to purchaser in a form suitable for application to plants, trees and other agricultural crops and hereby further represents and warrants that the fertilizer product shall meet the guaranteed minimum analysis as set forth on the product's identity label. However, this warranty does not extend to the use of this product contrary to label instructions and that the buyer assumes the risk of any such use.

Avoid breathing dust and prolonged exposure to dermis. Do not apply when weather condition favors drift from the areas treated. Do not apply when runoff is likely to occur. Do not use in a manner or at a time other than in a accordance with label directions because animal, plant or crop injury, illegal residues or other undesirable results may occur.

WARNING: Application of fertilizing materials containing Molybdenum may result in forage crops containing levels of Molybdenum which are toxic to ruminant animals. This product also contain Boron and may result in injury to crops.

NOTE: Information regarding the contents and levels of metals in this product is available on the Internet at http://www.aapfca.org/metals.htm.

