



FIFRA Section 2(ee) Recommendation

Date: July 14, 2020

Product: METOLACHLOR 8E (EPA Reg. No. 42750-340)

Use: Soybeans: postemergence (From emergence up through the 5th trifoliolate leaf stage).

States: AR, CO, FL, GA, IA, IL, IN, KS, KY, LA, MI, MN, MO, MS, NC, ND, NE, NY, OH, OK, PA, SC, SD, TN, TX, VA, WI, WV

DIRECTIONS FOR USE

For postemergence treatments, apply 1.0 to 1.33 pints per acre of METOLACHLOR 8E to soybeans from emergence through the 5th trifoliolate leaf stage. This product will not control emerged weeds, it must be applied to a weed free surface or in tank mixture with products that provide postemergence control of weeds present at the time of application. USE RESTRICTION: Make post emergence applications at least 90 days before harvest. USE PRECAUTION: Crop response can occur when postemergence applications are made during times of rapid growth and prolonged sunshine.

The user must refer to the federally approved METOLACHLOR 8E label and read and follow all applicable directions for use, restrictions, and precautions.

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This labeling must be in the possession of the user at the time of pesticide application.

This recommendation should be in the possession of the user at the time of pesticide application.

Use of this product according to this bulletin has not been reviewed or endorsed by the Office of the Indiana State Chemist.

This recommendation is made as permitted under FIFRA Section 2(ee), as amended, and has not been submitted to or accepted by the EPA.