



# Supplemental Label

## MOCCASIN® II PLUS Herbicide

(EPA Reg. No. 70506-333)

### FOR USE ON SUGARCANE

#### WARNING

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This supplemental label expires July 26, 2024 and must not be used or distributed after this date.

Read the label affixed to the container for MOCCASIN II PLUS Herbicide before applying. Use of MOCCASIN II PLUS Herbicide according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for MOCCASIN II PLUS Herbicide. This supplemental label and the EPA approved label must be in the possession of the user at the time of pesticide application.

Apply MOCCASIN II PLUS Herbicide as a preplant, preemergence, or postemergence treatment for weed control in sugarcane. It may also be used in a treatment program that includes a preplant/preemergence application followed by a postemergence/post-directed application.

#### Restrictions:

- **DO NOT** apply more than 1.71 qts/A (3.26 lbs ai) of MOCCASIN II PLUS Herbicide per year.
- **DO NOT** apply more than 1.22 qts/A (2.34 lbs ai) of MOCCASIN II PLUS Herbicide as a preplant or preemergence application per year.
- **DO NOT** apply more than 0.98 qts/A (1.88 lbs ai) of MOCCASIN II PLUS Herbicide as a postemergence application per year.
- **DO NOT** apply more than 0.49 qts/A (0.94 lbs ai) of MOCCASIN II PLUS Herbicide as a postemergence application if a preplant or preemergence application was made.
- **DO NOT** make more than two applications per year.
- **DO NOT** make applications less than 14 days apart.
- **DO NOT** apply to sugarcane that is taller than 60 inches.
- **DO NOT** make a postemergence application within 100 days of harvest.
- **DO NOT** exceed the total combined maximum annual sugarcane rates for s-metolachlor container products.

Apply MOCCASIN II PLUS Herbicide at a rate of 0.89 – 1.22 qts/A before planting, preemergence after new plantings, or after harvest, but before re-emergence of ratoon-cane. Use the higher rate on heavier soils and soils with higher organic matter content.

### **Postemergence Applications**

MOCCASIN II PLUS Herbicide will not control emerged weeds so apply only to a weed-free soil surface or in tank mixture with products that provide postemergence control of weeds present at the time of application. Apply MOCCASIN II PLUS Herbicide at 0.49 – 0.98 qts/A postemergence before the sugarcane reaches 60 inches in height. Postemergence applications may be made as broadcast post-over-the-top or as post-directed spray to soil between the rows and the base of the sugarcane.

If a preemergence application was made earlier in the season (not to exceed 1.22 qts/A), only 0.49 qts/A may be applied postemergence. The total amount of MOCCASIN II PLUS Herbicide applied (preemergence + postemergence) may not exceed 1.71 qts/A/year.

### **SUGARCANE - MOCCASIN II PLUS Herbicide TANK MIX COMBINATIONS**

It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

This product may be tank mixed with other registered sugarcane herbicides, insecticides, and fungicides. The tank mix partners are to be applied by the same methods and the same timings as MOCCASIN II PLUS Herbicide unless otherwise specified in the tank mix product label. Perform a compatibility test before spraying the tank mix application.

Rev. 7/22/21