

SUPPLEMENTAL LABELING

FENAZAQUIN	GROUP	21A	ACARICIDE
FENAZAQUIN	GROUP	39	FUNGICIDE

MAGISTER SC MITICIDE

EPA REG. No. 10163-322

TO ALLOW FOR AERIAL APPLICATIONS ON TREE NUTS AND STONE FRUIT
This supplemental label expires on October 5, 2024 and must not be used or distributed after this date.

ACTIVE INGREDIENT: Fenazaquin

4-tert-butylphenethyl quinazolin-4-yl ether (IUPAC).....18.79%

OTHER INGREDIENTS:.....81.21%

TOTAL 100.0%

Contains 1.7 pounds active ingredient per gallon.

KEEP OUT OF REACH OF CHILDREN WARNING/AVISO

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of pesticide application.
- Read the label affixed to the container for Magister SC Miticide before applying. Carefully following all precautionary statements and applicable use directions
- Use of Magister SC Miticide according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Magister SC Miticide.

DIRECTIONS FOR USE


RESTRICTIONS ON ALL CROPS:

- Do not apply by any type of irrigation system.
- Only ground sprayer, airblast and aerial applications are permitted.
- Do not apply during conditions when product will drift to other crops and non-target areas.
- Do not apply within 100 feet of water bodies (including, but not limited to, lakes; reservoirs; rivers; permanent streams; marshes or natural ponds; estuaries; and commercial fish farm ponds).
- Do not plant rotational crops, other than those on this label, within 30 days of this application. Do not plant root, tuber and bulb vegetables within 120 days of this application.

The required days between the last application and harvest are given in () after each crop name.

APPLICATION DIRECTIONS

CROP	PEST	RATE FL OZ/ACRE	LB A.I./ACRE	INSTRUCTIONS
STONE FRUIT GROUP 12-12 (3) Apricot; apricot, Japanese; capulin; cherry, black; cherry, Nanking; cherry, sweet; cherry, tart; Jujube, Chinese; nectarine; peach; plum; plum, American; plum, beach; plum, Canada; plum, cherry; plum, Chickasaw; plum, Damson; plum, Japanese; plum, Klamath; plum, prune; plumcot; sloe; cultivars, varieties, and/or hybrids of these	Mites, Powdery Mildew, Alternaria Leaf Spot* * suppression only	24 - 36	0.32 – 0.48	Ground: Apply in at least 50 gallons of water per acre. Use higher rates for heavier mite pressure. Air Application: Apply the recommended rate of Magister SC by air in a minimum of 10 GPA. Applications made by air to dense foliage may not provide adequate coverage of lower leaf surfaces for sufficient control. Use of higher labeled rates may be necessary.
Restrictions: <ul style="list-style-type: none">• Do not make more than one pre-harvest application per year.• Do not make more than one post-harvest application per year.• Do not exceed application rate of 36 fl oz of product (0.48 lb a.i.) per acre per year before harvest.• Do not exceed application rate of 36 fl oz of product (0.48 lb a.i.) per acre per year after harvest.• Do not exceed a maximum annual application rate of 0.96 lb a.i. per acre per year.• Pre-harvest: Do not apply this product until crop is through petal fall.				

<p>TREE NUT CROP GROUP 14-12 (7)</p> <p>African nut-tree; almond; beechnut; Brazil nut; Brazilian pine; bunya; bur oak; butternut; Cajou nut; candlenut; cashew; chestnut; chinquapin; coconut; coquito nut; dika nut; ginkgo; Guiana chestnut; hazelnut (filbert); heartnut; hickory nut; Japanese horse- chestnut; macadamia nut; mongongo nut; monkey-pot; monkey puzzle nut; Okari nut; Pachira nut; peach palm nut; pecan; pequi; Pili nut; pine nut; pistachio; Sapucaia nut; tropical almond; walnut, black; walnut, English; yellowhorn; cultivars, varieties, and/or hybrids of these</p>	 <p>Mites, Powdery Mildew, Shot Hole*, Rust*, Scab*, Alternaria Leaf Spot (suppression only)</p> <p><i>* suppression in almonds only</i></p>	<p>24 - 36</p>	<p>0.32 – 0.48</p>	<p>Ground: Apply in at least 100 gallons of water per acre. Slower application speed helps ensure good coverage. Use higher rates for heavier mite pressure.</p> <p>Air Application: Apply the recommended rate of Magister SC by air in a minimum of 10 GPA. Applications made by air to dense foliage may not provide adequate coverage of lower leaf surfaces for sufficient control. Use of higher labeled rates may be necessary.</p>
<p>Restrictions:</p> <ul style="list-style-type: none"> • Do not make more than one application per year. • Do not exceed application rate of 36 fl oz of product (0.48 lb a.i.) per acre per year. • Do not apply this product until crop is through petal fall. 				