

Maximum N-PACT

12 - 0 - 0

GUARANTEED ANALYSIS

Derived from: Urea-Triazone Solution, Boric Acid

*3.96% Slowly Available Nitrogen from Urea-Triazone Solution

WARNING: This product contains boron (B) and may be harmful to certain crops. Use only according to manufacturer's directions.

F1548

KEEP OUT OF REACH OF CHILDREN



May be harmful if swallowed. Causes skin irritation. Causes eye irritation. May cause respiratory irritation.

Density: 10.51 lbs/Gallon @ 68 °F

Net Contents: 2.50 Gallons (9.46 liters)

Net Weight: 26.28 lbs (11.92 kg)



MaxN-PactB C2719

FIRST AID

If in eyes: Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If on skin: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.

If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

Finhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice

FOR A MEDICAL EMERGENCY INVOLVING THIS PRODUCT, CALL 1-866-944-8565.

FOR FOLIAR / SOIL APPLICATION

DIRECTIONS FOR USE

Apply as needed based on soil tests or according to recommendations in your nutrient management plan.

APPLICATION INSTRUCTIONS

CROPS	FOLIER APPLIED RATE	APPLICATION TIMING	COMMENTS
Vegatable, Root and Tuber (01) Potato, Sugarbeet etc	1.0 to 6.0 pts/A	Apply prior to row canopy closing	
Leafy Veg (04) Celery Lettuce (Head, Leaf) etc	1.0 to 6.0 pts/A	Apply starting at thinning	
Brassica Veg (05) Broccoli, Cabbage, Cauliflower, Collards, Brussels Sprouts etc	1.0 to 6.0 pts/A	Apply at bottom establishment	
Legume Veg (06) All beans, Soybeans, etc	1.0 to 8.0 pts/A	Apply at early bloom	
Vegetable, Fruiting (08) Peppers, Tomato, Eggplant etc	1.0 to 6.0 pts/A	Apply starting at fruit formation	
Cucurbits (09) All melons, Cucumber, Squash etc	1.0 to 6.0 pts/A	Apply at bud formation to fruit establishment	
Citrus Fruits (10) Oranges, Lemons, Limes, etc	1.0 to 6.0 pts/A	Apply starting at fruit set	
Pome Fruit (11) Apple, Pear, etc	1.0 to 6.0 pts/A	Apply starting at fruit set	
Stone Fruit (12) Peach, Plum, etc	1.0 to 6.0 pts/A	Apply starting at fruit set	
Berry and Small Fruit (13) Strawberry, Blueberry, Caneberries (Blackberry, raspberry) etc	1.0 to 6.0 pts/A	Apply starting at fruit set	
Berry and Small Fruit (13) Grape	1.0 to 6.0 pts/A	Apply starting at fruit set through verasion	
Treenuts (14) Almond, Filbert, Pistachio, Pecan, Walnut etc	1.0 to 6.0 pts/A		
Cereal Grains (15) Corn, Wheat, Grain Sorghum, Rice, etc	1.0 to 6.0 pts/A	Apply from V5 to VT growth stage	
Grass forage Fodder and Hay (16) Grasses, etc.	1.0 to 6.0 pts/A		
Grass forage Fodder and Hay (17) Grass Grown for Seed	1.0 to 4.0 pts/A		
Animal Feed, Non Grass (18) All Alfalfa, Clover, etc	1.0 to 6.0 pts/A	Apply at crown green up or on regrowth after cutting	
Oilseed (20) Cotton, Sunflower, Canola etc	1.0 to 6.0 pts/A	Apply at bud to bolting stage	
Misc (99) Avocado, Artichoke, Hops, Flax, Mint, Olive, Peanut etc	1.0 to 4.0 pts/A		
Misc (99) Tobacco*	1.0 to 4.0 pts/A	Apply just prior to tasseling	*Do not use with alcohol based sucker control products

Crop Safety

This product has been tested on a number of crop varieties with no phytotoxicity observed at labeled rates; however, this product has not been tested on all varieties and cultivars. Due to the large number of plant species and their associated varieties or cultivars, and due to variable growing conditions, it is impossible to test every plant for tolerance to this product. It is the responsibility of the user to determine if this product can be used safely prior to commercial use. In a small test area, test the specified rates on plants for phytotoxicity prior to general use. Before using any tank mix, test the combination on a small portion of the crop to be treated to ensure that a phytotoxic response will not occur as a result of application. Treat a small group of target crop including plant cultivars and evaluate for 3 to 7 days prior to making an application to your entire crop.

Tank Mix Compatibility

This product is compatible with many agricultural pesticides. However, if specific information is not available, a jar test for compatibility should be conducted prior to large scale mixing.

STORAGE AND DISPOSAL

STORAGE: Store in original containers only. Keep containers tightly closed when not in use. Store in a cool, dry well-ventilated area, preferably in a locked storage area away from children, feed and food products, and seed. Store at ambient temperatures. Do not contaminate water, food, or feed by storage or disposal.

DISPOSAL: Do not reuse container. Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning, If burned, stay out of smoke. Do not contaminate water, food, or feed by storage or disposal.

HANDLING: Avoid breathing fertilizer dusts. Avoid prolonged or repeated skin contact. Avoid eve contact. Wash hands thoroughly after handling this product.

For help with any spill, leak, fire or exposure involving this material, call day or night CHEMTREC 1-800-424-9300.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

BEFORE BUYING OR USING THIS PRODUCT, read the Directions for Use and the following Conditions of Sale and Limitation of Warranty and Limitation of Warranty

Information regarding the contents and levels of metals in this product is available on the Internet at http://www.regulatorv-info-lpi.com

Tests have not been carried out under all possible use conditions and Seller cannot and does not represent or warrant that the goods are compatible with all other chemicals or under all use conditions.

Maximum N-Pact is a registered trademark of Loveland Products. Inc.



WARNING

This product can expose you to chemicals including nickel, which are known to the State of California to cause cancer. For more information go to www.P65Warnings.ca.gov.

GUARANTEED ANALYSIS

Total Nitrogen (N))%
8.04% Ŭrea Nitrogen	
3.96% Other Water Soluble Nitrogen*	
Boron (B)	ን%
5551 (5)	,,,

Derived from: Urea-Triazone Solution, Boric Acid

*3.96% Slowly Available Nitrogen from Urea-Triazone Solution

WARNING: This product contains boron (B) and may be harmful to certain crops. Use only according to manufacturer's directions.

KEEP OUT OF REACH OF CHILDREN



May be harmful if swallowed. Causes skin irritation. Causes eye irritation. May cause respiratory irritation.

