



Bayer Environmental Science  
A Division of Bayer CropScience LP  
5000 CentreGreen Way, Suite 400  
Cary, NC 27513  
1-866-99BAYER (1-866-992-2937)

## METHOD 240SL HERBICIDE®

For the tank mix application of Method 240SL Herbicide plus triclopyr amine (Element 3A or Garlon 3A) for control of mesquite and/or huisache in non-crop sites.

EPA Reg. No.  
Method 240SL: 432-1565  
Element 3A: 62719-37  
Garlon 3A: 62719-37

### **FIFRA 2(ee) Recommendation**

#### **FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF TEXAS**

This recommendation is valid until December 31, 2026

**FIFRA Section 2(ee) Recommendation:** This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. This product bulletin should be in the possession of the user at the time of application of the product. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

#### **DIRECTIONS FOR USE:**

It is a violation of federal law to use this product in a manner inconsistent with its label.

METHOD® 240SL herbicide can be tank mixed with triclopyr amine for control of mesquite and/or huisache in labeled non-crop sites. Apply METHOD® 240SL herbicide at rates of 8 to 12 fluid ounces product per acre in a tankmix with triclopyr at the rate of 4 to 6 ounces active ingredient per acre.

For mesquite, apply in the late-spring to mid-summer. For huisache, apply from late-summer into fall. Make applications when foliage is healthy, mature with a dark green color and when soil temperature is at least 75 degrees F at a depth of 12 inches. Control results are best when applications are made with moderate to high soil moisture and with uniform healthy foliage. Use an MSO adjuvant at 0.5% to 1% v/v in broadcast applications.

For best results with aerial applications, use at least 4 gpa carrier volume and apply using nozzles that deliver coarse or smaller droplets (VMD <450 microns) as defined by ASABE S572 standard.

For best results on huisache, use the high end of the rate range.

Before using Bayer METHOD® 240SL and triclopyr amine read and follow all applicable directions, restrictions and precautions on the EPA-registered label for each product.

#### **IMPORTANT PRECAUTIONS:**

Certain species may, in particular, be sensitive to low levels of METHOD® 240SL including but not limited to, conifers (such as Douglas fir, Norway spruce, ponderosa pine and white pine), deciduous trees (such as aspen, Chinese tallow, cottonwood, honey locust, magnolia, poplar species, redbud, silver maple, and willow species) and ornamental shrubs (such as arborvitae, burning bush, crape myrtle, forsythia, hydrangea, ice plant, magnolia, purple plum and yew).

Injury or loss of desirable trees or vegetation may result if METHOD® 240SL is applied on or near desirable trees or vegetation, on areas where their roots extend, or in locations where the treated soil may be washed or moved into contact with their roots. Consider site-specific characteristics and conditions that could contribute to unintended root zone exposure to desirable trees or vegetation. Root zone areas of desirable trees or vegetation are affected by local conditions and can extend beyond the tree canopy. If further information is needed regarding root zone area, consult appropriate state extension service, professional consultant or other qualified authority



Bayer Environmental Science  
A Division of Bayer CropScience LP  
5000 CentreGreen Way, Suite 400  
Cary, NC 27513  
1-866-99BAYER (1-866-992-2937)

## METHOD 240SL HERBICIDE®

For the tank mix application of Method 240SL Herbicide plus triclopyr amine (Element 3A or Garlon 3A) for control of mesquite and/or huisache in non-crop sites.

EPA Reg. No.  
Method 240SL: 432-1565  
Element 3A: 62719-37  
Garlon 3A: 62719-37

Injury to or loss of desirable trees or vegetation may result if equipment is drained or flushed on or near these trees or vegetation, or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots.

### IMPORTANT RESTRICTIONS:

DO NOT USE PLANT MATERIAL TREATED WITH METHOD® 240SL FOR MULCH OR COMPOST.

Do not apply this product in areas where the roots of desirable trees and/or shrubs may extend unless injury or loss can be tolerated. Root zone areas of desirable trees or vegetation are affected by local conditions and can extend well beyond the tree canopy.

Do not apply this product if site-specific characteristics and conditions exist that could contribute to movement and unintended root zone exposure to desirable trees or vegetation unless injury or loss can be tolerated.

Do not apply METHOD 240SL HERBICIDE to highways/roadsides or other non-crop areas during periods of intense rainfall or where prevailing soils are either saturated with water or of a type through which rainfall will not readily penetrate, as this may result in off-site movement.

Do not apply more than 18 fluid ounces METHOD® 240SL herbicide per acre per year on a broadcast acre basis.

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577.  
For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937).

*As with any herbicide product, always read and follow label directions.  
For additional information call toll-free 1-866-99BAYER (1-866-992-2937).*