
SUPPLEMENTAL LABELING

Syngenta Crop Protection, LLC

P. O. Box 18300

Greensboro, North Carolina 27419-8300

SCP 1603A-S3 0223

PYDIFLUMETOFEN	GROUP	7	FUNGICIDE
FLUDIOXONIL	GROUP	12	FUNGICIDE

Miravis® Prime

Fungicide

ADEPIDYN® Technology*

This supplemental label expires on 08/23/2024 and must not be used or distributed after this date.

Active Ingredients:

Pydiflumetofen**: 12.8%

Fludioxonil***: 21.4%

Other Ingredients: 65.8%

Total: 100.0%

*Technology denotes the active ingredient Pydiflumetofen

**CAS No. 1228284-64-7

***CAS No. 131341-86-1

Miravis® Prime is formulated as a suspension concentrate and contains 1.25 lb of pydiflumetofen and 2.09 lb fludioxonil per gallon.

KEEP OUT OF REACH OF CHILDREN.**CAUTION**

See additional precautionary statements and directions for use inside booklet. See First Aid Statement inside booklet and on container label.

EPA Reg. No. 100-1603

All applicable directions, restrictions, and precautions on the EPA-registered label are to be followed. Before using Miravis Prime as permitted according to this supplemental label, read and follow all applicable directions, restrictions, and precautions on the EPA-registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and/or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.



DIRECTIONS FOR USE

CROP USE DIRECTIONS

Caneberry Crop Subgroup 13-07A

Crops (Including all cultivars, varieties and/or hybrids of these)			
Blackberry Loganberry Raspberry, red and black Wild raspberry			
Target Disease	Rate (fl oz/A)	Application Timing	Use Directions
Alternaria fruit rot (<i>Alternaria</i> spp.) Powdery mildew (<i>Sphaerotheca macularis</i>) Spur blight (<i>Didymella</i> spp.)	9 – 13.4*	Begin applications prior to disease development. Continue applications on a 7-day interval, following the resistance management guidelines.	Apply by ground or air. An adjuvant may be added at specified rates. Apply in sufficient water volume to ensure good coverage.
Botrytis fruit rot (<i>Botrytis cinerea</i>)	10.3 – 13.4*		If disease pressure is high, use the highest rate.
*9.0 fl oz product/A is equivalent to 0.088 lb ai pydiflumetofen and 0.147 lb ai fludioxonil. *13.4 fl oz product/A is equivalent to 0.131 lb ai pydiflumetofen and 0.219 lb ai fludioxonil.			
Resistance Management: • Do not make more than two consecutive applications of Miravis Prime or other Group 7 and 12 fungicides before alternation with a fungicide that is not in Group 7 or 12.			
USE RESTRICTIONS			
1) Maximum Single Application Rate: DO NOT exceed the maximum rate listed in the table. 2) Maximum Number of Applications per Year: DO NOT make more than 2 applications at the maximum application rate per year. 3) Minimum Application Interval: 7 days 4) Maximum Annual Rate: 26.8 fl oz/A/year (equivalent to 0.262 lb ai pydiflumetofen and 0.438 lb ai fludioxonil) a. DO NOT apply more than 0.268 lb ai/A/year of pydiflumetofen-containing products. b. DO NOT apply more than 0.9 lb ai/A/year of fludioxonil-containing products. 5) Pre-harvest Interval (PHI): 0 days 6) Make no more than two applications by air per year.			

Peppers, Greenhouse Production Only

Crops (Including all cultivars, varieties, and/or hybrids of these)			
Bell pepper Nonbell pepper			
Target Disease	Rate (fl oz/A)	Application Timing	Use Directions
Suppression Only: Fusarium wilt (<i>Fusarium</i> spp.)	11.4*	Begin applications prior to disease development.	Apply via drench or drip irrigation. See Section 4.1.1 and Section 4.5.4 for directions on drip irrigation application.
*11.4 fl oz product/A is equivalent to 0.11 lb ai pydiflumetofen and 0.186 lb ai fludioxonil.			
USE RESTRICTIONS			
<ol style="list-style-type: none"> 1) Maximum Single Application Rate: Do not exceed the maximum rate listed in the table. 2) Maximum Number of Applications per Year: Do not make more than 1 soil application at the maximum application rate per year. 3) Minimum Application Interval: Not Applicable 4) Maximum Annual Rate: 40.2 fl oz/A/year (equivalent to 0.393 lb ai/A/year pydiflumetofen and 0.654 lb ai/A/year fludioxonil). <ol style="list-style-type: none"> a. Do not apply more than 0.393 lb ai/A/year of pydiflumetofen-containing products. b. Do not apply more than 0.9 lb ai/A/year of fludioxonil-containing products. 5) Do not apply to fruiting vegetables grown in the greenhouse except for greenhouse peppers. 6) Pre-harvest Interval (PHI): <ol style="list-style-type: none"> a. Soil application: 1 day 			

ADEPIDYN®, Miravis® and the Syngenta logo are trademarks of a Syngenta Group Company
©2023 Syngenta