

---

**SUPPLEMENTAL LABELING**

---

**Syngenta Crop Protection, LLC**  
P. O. Box 18300  
Greensboro, North Carolina 27419-8300

**SCP 1601A-S2 0523**

<b>PYDIFLUMETOFEN</b>	<b>GROUP</b>	<b>7</b>	<b>FUNGICIDE</b>
-----------------------	--------------	----------	------------------

**Miravis**

Fungicide

ADEPIDYN® Technology\*

**This supplemental label expires on 11/18/2024 and must not be used or distributed after this date.**

Active Ingredient(s):	
Pydiflumetofen** .....	18.3%
Other Ingredients:	81.7%
Total:	100.0%

\*Technology denotes the active ingredient Pydiflumetofen.

\*\*CAS No. 1228284-64-7

Miravis is formulated as a suspension concentrate (SC) and contains 1.67 lb of pydiflumetofen per gallon.

**KEEP OUT OF REACH OF CHILDREN.**

**CAUTION**

**EPA Reg. No. 100-1601**

**All applicable directions, restrictions, and precautions on the EPA-registered label are to be followed. Before using Miravis as permitted according to this supplemental label, read and follow all applicable directions, restrictions, and precautions on the EPA-registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and/or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.**

**syngenta.**

## DIRECTIONS FOR USE

### CROP USE DIRECTIONS

#### Lettuce, Greenhouse Production Only

Crops (Including all cultivars and/or varieties of these) Not for use in California			
Lettuce, head Lettuce, leaf			
Target Disease	Rate fl oz/A (lb ai/A)	Application Timing	Use Directions
<b>Foliar Application:</b> Powdery mildew <i>(Erysiphe cichoracearum)</i> Gray mold <i>(Botrytis cinerea)</i>	5.1 (0.067)	Begin applications prior to disease development.  Continue applications on a 7-day interval, following resistance management guidelines.	Apply to foliage.  An adjuvant may be added at specified rates.
<b>Resistance Management:</b> <ul style="list-style-type: none"> <li>Refer to <b>Section 3.2</b>.</li> <li>Do not make more than two consecutive applications of Miravis or other Group 7 fungicides before alternation with a fungicide that is not in Group 7.</li> </ul>			
USE RESTRICTIONS			
1) Refer to <b>Section 6.1</b> for additional product use restrictions. 2) <b>Maximum Single Application Rate: DO NOT</b> exceed the maximum rate listed in the table. 3) <b>Maximum Number of Applications per Year: DO NOT</b> make more than 2 foliar applications at the maximum application rate per year. 4) <b>Minimum Application Interval:</b> 7 days 5) <b>Maximum Annual Rate:</b> 27.4 fl oz/A/year (equivalent to 0.357 lb ai/A/year of pydiflumetofen-containing products). 6) <b>DO NOT apply to leafy green vegetables grown in the greenhouse except for greenhouse lettuce.</b> 7) <b>Pre-harvest Interval (PHI):</b> 0 days			

Miravis®, Adepidyn® and the Syngenta logo are trademarks of a Syngenta Group Company

©2023 Syngenta