
SUPPLEMENTAL LABELING

Syngenta Crop Protection, LLC
P. O. Box 18300
Greensboro, North Carolina 27419-8300

SCP 1602A S1 0819

DIFENCONAZOLE	GROUP	3	FUNGICIDE
PYDIFLUMETOFEN	GROUP	7	FUNGICIDE

Miravis® Top

Fungicide

This supplemental label expires on 08/30/2022 and must not be used or distributed after this date.

Active Ingredients:

Pydiflumetofen*:	6.9%
Difenoconazole**:	11.5%
Other Ingredients:	81.6%
Total:	100.0%

*CAS No. 1228284-64-7

**CAS No. 119446-68-3

MIRAVIS TOP is formulated as a suspension concentrate and contains 0.63 lb of pydiflumetofen and 1.04 lb difenoconazole per gallon.

KEEP OUT OF REACH OF CHILDREN.

CAUTION

EPA Reg. No. 100-1602

All applicable directions, restrictions, and precautions on the EPA-registered label are to be followed. Before using MIRAVIS TOP as permitted according to this supplemental label, read and follow all applicable directions, restrictions, and precautions on the EPA-registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and/or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

syngenta.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

CROP USE DIRECTIONS

Citrus Fruit, Crop Group 10-10

Crops (Including cultivars, varieties, and/or hybrids of these)						
Australian desert lime	Lemon	Satsuma mandarin				
Australian finger-lime	Lime	Sweet lime				
Australian round lime	Mediterranean mandarin	Tachibana orange				
Brown River finger lime	Mount white lime	Tahiti lime				
Calamondin	New Guinea wild lime	Tangelo				
Citron	Orange, sour	Tangerine (mandarin)				
Citrus hybrids	Orange, sweet	Tangor				
Grapefruit	Pummelo	Trifoliolate orange				
Japanese summer grapefruit	Russell River lime	Uniq fruit				
Kumquat						
Target Disease	Rate (fl oz/A)	Application Timing	Use Directions			
Albinism (<i>Alternaria alternata</i> pv <i>citri</i>)	13.7 - 15.2*	Begin applications prior to disease development. Continue throughout the season on 7- to 21-day interval following the resistance management guidelines.	Apply by ground or air. An adjuvant may be added at recommended rates. Apply in sufficient water to obtain thorough coverage. A horticultural spray oil should be used to improve control of greasy spot. If disease pressure is high, use the shortest interval and highest rate.			
Alternaria Leaf and Fruit Spot (<i>Alternaria citri</i>)						
Black Spot (<i>Guignardia citricarpa</i>)						
Diplodia Stem-End Rot (<i>Diplodia natalensis</i>)						
Melanose (<i>Diaporthe citri</i>)						
Phomopsis Stem-End Rot (<i>Phomopsis citri</i>)						
Scab (<i>Elsinoe fawcettii</i>)						
Sweet Orange Scab (<i>Elsinoe australis</i>)						
Anthracnose (<i>Colletotrichum</i> spp.)				15.2*		
Post bloom fruit drop (PFD) (<i>Colletotrichum acutatum</i>)						
*13.7 fl oz product/A is equivalent to 0.067 lb ai pydiflumetofen and 0.111 lb ai difenoconazole. *15.2 fl oz product/A is equivalent to 0.075 lb ai pydiflumetofen and 0.124 lb ai difenoconazole.						
Resistance Management:						
<ul style="list-style-type: none"> • Do not make more than two consecutive applications of Miravis Top or other Group 3 and 7 fungicides before alternation with a fungicide that is not in Group 3 or 7. 						
USE RESTRICTIONS						

- 1) Refer to **Section 6.1** for additional product use restrictions.
- 2) **Maximum Single Application Rate:** Do not exceed the maximum rate listed in the table.
- 3) **Maximum Number of Applications per Year:** Do not make more than 4 applications at the maximum application rate per year.
- 4) **Minimum Application Interval:** 7 days
- 5) **Maximum Annual Rate:** 60.9 fl oz/A/year (equivalent to 0.30 lb ai pydiflumetofen and 0.496 lb ai Difenconazole)
 - a. **Do not** apply more than 0.50 lb ai/A/year of difenoconazole-containing products.
 - b. **Do not** apply more than 0.30 lb ai/A/year of pydiflumetofen-containing products.
- 6) **Pre-harvest Interval (PHI):** 0 days
- 7) For aerial applications do not apply in less than 10 GPA water.

Cotton (Cottonseed Subgroup 20C)

Crops (Including cultivars, varieties, and/or hybrids of these)			
Cottonseed			
Target Disease	Rate (fl oz/A)	Application Timing	Use Directions
Alternaria leafspot (<i>Alternaria</i> spp.)	13.6*	Begin applications prior to disease development.	Apply by ground, air, or chemigation.
Areolate mildew (<i>Ramularia gossypii</i>)		Continue applications through season on a 14-day interval, following the resistance management guidelines.	An adjuvant may be added at recommended rates.
Ascochyta Blight (<i>A. gossypii</i>)			Apply in sufficient water to obtain thorough coverage.
Boll Rots (<i>Ascochyta gossypii</i> , <i>Alternaria</i> spp., <i>Diplodia</i> spp., <i>Phoma</i> spp.)		For Target Spot: The first application should begin at first bloom. Depending on environmental conditions and the health of the cotton plant, additional applications may be made on a 14 to 21 day interval.	If disease pressure is high, use the highest rate.
Hardlock (<i>Fusarium verticillioides</i>)			
Leafspots and blights (<i>Alternaria</i> spp., <i>Ascochyta gossypii</i> , <i>Cercospora</i> spp., <i>Stemphyllium</i> spp.)			
Target spot (<i>Corynespora cassiicola</i>)			
*13.6 fl oz product/A is equivalent to 0.067 lb ai pydiflumetofen and 0.111 lb ai Difenconazole.			
Resistance Management:			
<ul style="list-style-type: none"> • Do not make more than two consecutive applications of Miravis Top or other Group 3 and 7 fungicides before alternation with a fungicide that is not in Group 3 or 7. 			
USE RESTRICTIONS			
<ol style="list-style-type: none"> 1) Refer to Section 6.1 for additional product use restrictions. 2) Maximum Single Application Rate: Do not exceed the maximum rate listed in the table. 3) Maximum Number of Applications per Year: Do not make more than 3 applications at the maximum application rate per year. 			

- 4) **Minimum Application Interval:** 14 days
- 5) **Maximum Annual Rate:** 40.8 fl oz/A/year (equivalent to 0.201 lb ai pydiflumetofen and 0.333 lb ai Difenoconazole)
 - a. **Do not** apply more than 0.34 lb ai/A/year of difenoconazole-containing products.
 - b. **Do not** apply more than 0.201 lb ai/A/year of pydiflumetofen-containing products.
- 6) **Pre-harvest Interval (PHI):** 45 days

Pecans

Crops (Including all cultivars, varieties, and/or hybrids of these)			
Pecans			
Target Disease	Rate (fl oz/A)	Application Timing	Use Directions
Downy Spot (<i>Mycosphaerella caryigena</i>)	13.6*	Begin applications prior to disease development.	Apply by ground or air.
Liver Spot (<i>Gnomonia caryae pv pecanae</i>)		Continue applications through season on a 14-day interval, following the resistance management guidelines.	An adjuvant may be added at recommended rates.
Pecan Scab (<i>Cladosporium caryigenum</i>)			If disease pressure is high, use the shortest interval.
Powdery Mildew (<i>Microsphaera penicillata</i>)			
Vein Spot (<i>Gnomonia nerviseda</i>)			
Zonate Leaf Spot (<i>Grovesinia pyramidalis</i>)			
*13.6 fl oz product/A is equivalent to 0.067 lb ai pydiflumetofen and 0.111 lb ai difenoconazole.			
Resistance Management:			
<ul style="list-style-type: none"> • Do not make more than two consecutive applications of Miravis Top or other Group 3 and 7 fungicides before alternation with a fungicide that is not in Group 3 or 7. 			
USE RESTRICTIONS			
<ol style="list-style-type: none"> 1) Refer to Section 6.1 for additional product use restrictions. 2) Maximum Single Application Rate: Do not exceed the maximum rate listed in the table. 3) Maximum Number of Applications per Year: Do not make more than 4 applications at the maximum application rate per year. 4) Minimum Application Interval: 14 days 5) Maximum Annual Rate: 54.4 fl oz/A/year (equivalent to 0.442 lb ai/A/year difenoconazole and 0.268 lb ai/A/year pydiflumetofen) <ol style="list-style-type: none"> a. Do not apply more than 0.46 lb ai/A/year of difenoconazole-containing products. b. Do not apply more than 0.268 lb ai/A/year of pydiflumetofen-containing products. 6) Pre-harvest Interval (PHI): Do not apply after shuck split or within 45 days of harvest. 7) For aerial applications do not apply in less than 10 GPA water. 			

Miravis® and the Syngenta logo are trademarks of a Syngenta Group Company

©2019 Syngenta

000100-01602.20180731C.MIRAVIS TOP_CROP_SUPP_JUL2018-CL.pdf