



# Supplemental Label

## MOCCASIN® II Plus Herbicide

(EPA Reg. No. 70506-333)

### FOR USE ON SOD FARMS

#### WARNING

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.  
This supplemental label expires July 26, 2024 and must not be used or distributed after this date.

Read the label affixed to the container for MOCCASIN II Plus Herbicide before applying. Use of MOCCASIN II Plus Herbicide according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for MOCCASIN II Plus Herbicide. This supplemental label and the EPA approved label must be in the possession of the user at the time of pesticide application.

**DO NOT USE IN GREENHOUSES OR OTHER ENCLOSED STRUCTURES.**

#### APPLICATION PROCEDURES

**Ground Application:** Apply MOCCASIN II Plus Herbicide alone or in tank mixtures by ground equipment in a minimum of 10 gallons of spray mixture per acre, unless otherwise specified.

Use sprayers that provide accurate and uniform application. For MOCCASIN II Plus Herbicide tank mixtures with wettable powder or dry flowable formulations, screens and strainers should be no finer than 50-mesh. Rinse sprayer thoroughly with clean water immediately after use.

Calculate the amount of herbicide needed for band treatment by the following formula:

$$\frac{\text{band width in inches}}{\text{row width in inches}} \times \text{broadcast rate per acre} = \text{amount needed per acre of field}$$

**Aerial Application (Sod Farms Only):** Apply MOCCASIN II Plus Herbicide in water alone or in tank mixtures with atrazine, simazine, or other herbicides registered for use on sod farms in a minimum total volume of 2 gal/A by aircraft. See Turfgrass section for a listing of applicable warm-season grasses. Avoid application under conditions where uniform coverage cannot be obtained or where excessive spray drift may occur. In order to assure that spray will be controllable within the target area when used according to label directions, apply at a maximum height of 10 ft, using low- drift nozzles at a maximum pressure of 40 psi, and restrict application to periods when wind speed is 10 mph or less. To assure that spray will not adversely affect adjacent sensitive nontarget plants, apply MOCCASIN II Plus Herbicide or MOCCASIN II Plus Herbicide mixtures at a minimum upwind distance of 400 ft from sensitive plants.

Avoid application to humans or animals. Flagmen and loaders must avoid inhalation of spray mist and prolonged contact with skin.

#### Turfgrass

**Warm Season Grasses (Bermudagrass, Centipedegrass, St. Augustinegrass, Bahiagrass, and Zoysiagrass) including Commercial St. Augustinegrass Sod Production**

Apply **MOCCASIN II Plus Herbicide before weeds emerge**. Since soil moisture is necessary for activation, irrigate with ½ inch of water if rainfall does not occur within 7 days after treatment (See following Precautions).

## Weeds Controlled

| Common Name         | Scientific Name                | Rate of MOCCASIN II Plus Herbicide* |
|---------------------|--------------------------------|-------------------------------------|
| Annual sedge        | <i>Cyperus compressus</i>      | 2.6 pts/A                           |
| Yellow nutsedge     | <i>Cyperus esculentus</i>      |                                     |
| Smooth crabgrass    | <i>Digitaria ischaemum</i>     |                                     |
| Large crabgrass     | <i>Digitaria sanguinalis</i>   |                                     |
| Bearded sprangletop | <i>Leptochloa fascicularis</i> | 1.3-2.6 pts/A                       |
| Mexican sprangletop | <i>Leptochloa uninervia</i>    |                                     |
| Doveweed            | <i>Murdannia nudiflora</i>     |                                     |
| Annual bluegrass    | <i>Poa annua</i>               |                                     |

\*1.0 pt/A = 0.3 ml/1,000 sq ft 1.3 pt/A = 0.4 ml/1,000 sq ft 2.6 pt/A = 0.9 ml/1,000 sq ft

## Restrictions:

**DO NOT** use MOCCASIN II Plus Herbicide on turfgrasses in New York State.

**DO NOT** use on golf greens, tees, or aprons

Split rate of applications can be made at rates not less than 1 pt/A (0.9 lbs ai/A). **DO NOT** apply more than 2.6 pts/A (2.4 lbs ai/A) in a single application.

**DO NOT** apply more than once every 6 weeks.

For commercial sod production, **DO NOT** apply more than 4.2 pts/A (4.0 lbs ai/A) per year to the same area used for sod production.

For commercial sod production, **DO NOT** make more than 4 applications per acre per year (not to exceed 4.2 pts/A (4.0 lbs ai/A) per year).

**DO NOT** graze or feed turf clippings to animals.

## Precautions for all uses on sod farms:

Delayed spring green-up, temporary slowing of growth and yellowing may occur following application. To avoid turf injury:

- Application of a nitrogen-containing fertilizer at or soon after applying MOCCASIN II Plus Herbicide will minimize delay in spring green-up and any temporary yellowing;
- Use only on turfgrass not under stress from infestations of insects, nematodes, or diseases;
- do not seed or overseed with desirable turfgrass 4 months before or after treatment, and
- do not apply this product to newly seeded grasses until they have overwintered and have a well-developed rhizome system.
- Before using MOCCASIN II Plus Herbicide in the tank mix with fluid fertilizer or other registered pesticides, determine the tolerance of the turf species by applying the combination to a limited area during a period of active growth.
- In turfgrass areas which have heavy thatch, the weed control of MOCCASIN II Plus Herbicide may be reduced.

Rev. 7/22/21