



Bayer CropScience LP
P.O. Box 12014
2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709
1-866-99BAYER (1-866-992-2937)

	GROUP	23	INSECTICIDE
MOVENTO®			
<i>EPA Reg. No. 264-1050</i>			
Control and Suppression of Pests Infesting Young Immature Tree Nut Crops Via Handgun Application			

FIFRA 2(ee) Recommendation

FOR DISTRIBUTION AND USE ONLY IN THE STATE OF CALIFORNIA

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or accepted by the U.S. Environmental Protection Agency. This product bulletin must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product inconsistent with its labeling.

TREE NUTS			
Crops of Crop Group 14 plus Pistachio Including: Almond, Beechnut, Brazil nut, Butternut, Cashew, Chestnut, Chinquapin, Filbert (hazelnut), Hickory nut, Macadamia nut (bush nut), Pecan, Pistachio, Walnut [black and English (Persian)]			
Pests Controlled		Product Rate	
Aphids Mealybugs Phylloxera San Jose scale Walnut scale Whiteflies	1st Year Trees		2nd Year Trees
	(fl oz/100 gallons)	(lb ai/100 gallons)	(fl oz/100 gallons)
	32.0	0.5	16.0
Pests Suppressed			
European fruit lecanium scale Nematodes Olive scale Pacific mite Twospotted spider mite		Apply in a minimum spray volume of 10 gallons per acre, maximum spray volume of 25 gallons per acre	Apply in a minimum spray volume of 20 gallons per acre, maximum spray volume of 50 gallons per acre
Foliar Application Restrictions:			
Pre-Harvest Interval (PHI): 7 day(s)			
Minimum interval between applications: 14 days			
Maximum MOVENTO allowed per application: 9 fluid ounces/Acre (0.14 lb AI/Acre)			
Maximum MOVENTO allowed per 12-month period: 21.5 fluid ounces/Acre (0.34 lb AI/Acre)			
Do not apply until after petal fall.			
Make application with a handgun sprayer to young immature reset or solid-set trees. MOVENTO must be tank-mixed with a spray adjuvant / additive having spreading and penetrating properties to maximize leaf uptake and systemicity of the active ingredient within treated plants; refer to adjuvant manufacturer's label for appropriate rates.			
For control of San Jose scale, apply immediately after petal fall; under heavy infestation pressure or where difficult control conditions exist, a second application may be necessary.			

For **MEDICAL** And **TRANSPORTATION** Emergencies **ONLY** Call 24 Hours A Day 1-800-334-7577

For **PRODUCT USE** Information Call 1-866-99BAYER (1-866-992-2937)

Visit our worldwide web site at www.bayercropscienceus.com

As with any crop-protection product, always read and follow label directions.

For additional information call toll-free 1-866-99BAYER (1-866-992-2937)