



Bayer CropScience

Bayer CropScience LP
P.O. Box 12014
2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709
1-866-99BAYER (1-866-992-2937)

	GROUP	23	INSECTICIDE
MOVENTO® MPC			
EPA Reg. No. 264-1065			
Use on Young Immature Citrus Trees - via Handgun or Hoopboom Application			

FIFRA 2(ee) Recommendation

FOR DISTRIBUTION AND USE ONLY IN THE STATES OF FLORIDA, ALABAMA OR GEORGIA

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. This product bulletin should be in the possession of the user at the time of application of the product. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its label.

CITRUS FRUITS

Crops of Crop Group 10-10 Including: Australian lime (desert, finger, and round), Brown River finger lime, Calamondin, Citron, Grapefruit, Japanese summer grapefruit, Kumquat, Lemon, Lime, Mediterranean mandarin, Mount White lime, New Guinea wild lime, Orange (sour and sweet), Pummelo, Russell River lime, Satsuma mandarin, Sweet lime, achibana orange, Tahiti lime, Tangelo, Tangerine, Tangor, Trifoliate orange, Uniq fruit, including cultivars, varieties and/or hybrids of these commodities.

PESTS CONTROLLED	Rate fluid ounces/100 gallons	Rate lb ai/100 gallons
Aphids Asian citrus psyllid California red scale Citrus leafminer Citrus bud mite Citrus red mite Citrus rust mite (silver mite) Citrus snow scale Citrus thrips Florida red scale Mealybugs Pink citrus rust mite Purple scale Texas citrus mite Whiteflies	24.0 – 32.0	0.23 – 0.31
PESTS SUPPRESSED		
Black scale Brown scale Citricola scale Cottony cushion scale Nematodes		

Notes and Restrictions

Pre-Harvest Interval (PHI): **1 day.**

Minimum interval between applications: **21 days.**

Maximum MOVENTO MPC allowed per 12 month period: **32 fluid ounces/Acre (0.32 lb ai/Acre).**

Do not apply more than 16 fluid ounces/Acre/Application (0.16 lb ai/Acre/Application).

Do not apply this product within 10 days prior to bloom, during bloom, or until petal fall is complete.

Do not apply non-ionic surfactants in tank mix combination with MOVENTO MPC on white grapefruit.

Make application with a handgun or hoopboom sprayer to young immature reset or solid-set citrus trees, utilizing sufficient spray volume for sufficient coverage of all plant parts.

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577.

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937).

As with any crop-protection product, always read and follow label directions.

For additional information call toll-free 1-866-99BAYER (1-866-992-2937).