



**Bayer CropScience**

**Bayer CropScience LP  
P.O. Box 12014  
2 T.W. Alexander Drive  
Research Triangle Park, North Carolina 27709  
1-866-99BAYER (1-866-992-2937)**

**MOVENTO®**

**EPA Reg. No. 264-1050**

For Use On: Bushberry; Carrot; Stone Fruits; Sugar Beet; Tree Nuts

*This supplemental label expires on 06/26/2020 and must not be used or distributed after this date.*

**Supplemental Label**

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation. Read this label and the product package label before using this product. This Supplemental Label must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the product label for MOVENTO® attached to the container.

**BUSHBERRY SUBGROUP  
LOW GROWING BERRY SUBGROUP**

**Crops of Crop Subgroups 13-07B and 13-07H Including:** Aronia berry, Bearberry, Bilberry, Blueberry (highbush and lowbush), Chilean guava, Cloudberry, Cranberry, Currant (black, buffalo, native, and red), Elderberry, European barberry, Gooseberry, Edible honeysuckle, Jostaberry, Juneberry, Muntries, Lingonberry, Partridgeberry, Salal, Sea buckthorn, and cultivars, varieties, and/or hybrids of these.

Pests Controlled	Product Rate	
	(fl oz/A)	(lb ai/A)
Aphids Blueberry Gall Midge Cranberry Tipworm Thrips (larvae)	8.0 - 10.0	0.13 - 0.16
Pests Suppressed	Product Rate	
	(fl oz/A)	(lb ai/A)
Blueberry Maggot Leafhoppers	10.0	0.16

**Foliar Application Restrictions:**  
Pre-Harvest Interval (PHI): **7 days**  
Minimum interval between applications: **7 days**  
Maximum MOVENTO allowed per calendar year: **30 fl oz/A**  
Maximum spirotetramat per crop season: **0.47 lb ai/A**  
Do not apply until after petal fall

**CARROT**

Pests Controlled	Product Rate	
	(fl oz/A)	(lb ai/A)
Aphids Whiteflies	5.0	0.08

**Foliar Application Restrictions:**  
Pre-Harvest Interval (PHI): **1 day**  
Minimum interval between applications: **7 days**  
Maximum MOVENTO allowed per crop season: **10 fl oz/A**  
Maximum spirotetramat per crop season: **0.16 lb ai/A**

*MOVENTO is a Registered trademark of Bayer.*

SAL 06/28/2017 – AV1

**STONE FRUITS**

**Crops of Crop Group 12-12 Including:** Apricot; apricot, Japanese; capulin; cherry, black; cherry, Nanking; cherry, sweet; cherry, tart; Jujube, Chinese; nectarine; peach; plum; plum, American; plum, beach; plum, Canada; plum, cherry; plum, Chickasaw; plum, Damson; plum, Japanese; plum, Klamath; plum, prune; plumcot; sloe; cultivars, varieties, and/or hybrids of these

Pests Controlled	Product Rate	
	(fl oz/A)	(lb ai/A)
Aphids Mealybugs San Jose scale White peach scale Whiteflies	6.0 – 9.0	0.09 - 0.14
<b>Pests Suppressed</b>		
Black scale Cherry fruit fly European fruit lecanium scale European red mite Nematodes Spotted Wing Drosophila Twospotted spider mite		

**Foliar Application Restrictions:**

Pre-Harvest Interval (PHI): **7 day(s)**

Minimum interval between applications: **14 days**

Maximum MOVENTO allowed per calendar year: **15.3 fl oz/A**

Maximum spirotetramat per calendar year: **0.24 lb ai/A**

Do not apply until after petal fall.

For control of San Jose scale, apply immediately after petal fall; under heavy infestation pressure or where difficult control conditions exist, a second application may be necessary.

**SUGAR BEET**

Pests Controlled	Product Rate	
	(fl oz/A)	(lb ai/A)
Bean aphid Root Aphid Whiteflies	5.0 - 9.0	0.08 - 0.14
<b>Pests Suppressed</b>		
Root Maggot Sugarbeet cyst nematode		

**Foliar Application Restrictions:**

Pre-Harvest Interval (PHI): **28 day(s)**

Minimum interval between applications: **14 days**

Maximum MOVENTO allowed per crop season: **18 fl oz/A**

Maximum spirotetramat per crop season: **0.28 lb ai/A**

**TREE NUTS**

**Crops of Crop Group 14-12 Including:** African nut-tree; almond; beechnut; Brazil nut; Brazilian pine; bunya; bur oak; butternut; Cajou nut; candlenut; cashew; chestnut; chinquapin; coconut; coquito nut; dika nut; ginkgo; Guiana chestnut; hazelnut (filbert); heartnut; hickory nut; Japanese horse-chestnut; macadamia nut; mongongo nut; monkey-pot; monkey puzzle nut; Okari nut; Pachira nut; peach palm nut; pecan; pequi; Pili nut; pine nut; pistachio; Sapucaia nut; tropical almond; walnut, black; walnut, English; yellowhorn; cultivars, varieties, and/or hybrids of these

Pests Controlled	Product Rate	
	(fl oz/A)	(lb ai/A)
Aphids Mealybugs Phylloxera San Jose scale Walnut scale Whiteflies	6.0 – 9.0	0.09 - 0.14
<b>Pests Suppressed</b>		
European fruit lecanium scale Nematodes Olive scale Pacific mite Twospotted spider mite		

**Foliar Application Restrictions:**

Pre-Harvest Interval (PHI): **7 day(s)**

Minimum interval between applications: **14 days**

Maximum MOVENTO allowed per calendar year: **21.5 fl oz/A**

Maximum spirotetramat per calendar year: **0.34 lb ai/A**

Do not apply until after petal fall.

For control of San Jose scale, apply immediately after petal fall; under heavy infestation pressure or where difficult control conditions exist, a second application may be necessary.

**For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577**

**For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)**

*As with any crop-protection product, always read and follow label directions.*

*For additional information call toll-free 1-866-99BAYER (1-866-992-2937).*