

**Muscle[®] 3.6 Fungicide****(EPA Reg. No. 60063-29)**

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

CROP	PHI (DAYS)	DISEASES	RATE OF PRODUCT PER ACRE	APPLICATION DIRECTIONS
Soybeans	21	Rust (<i>Phakopsora pachyrhizii</i>)	3 to 4 fl oz per acre Maximum 12 fl oz per acre per crop season	At the first sign of rust pustules on foliage make a spray application of this product. If environmental conditions are favorable to continued disease development, make a second application after 10 to 21 days dependent upon the severity of disease pressure. Do not apply more than 0.338 pounds of the active ingredient, tebuconazole, per acre per year and make no more than 3 applications of this product per season. Apply this product in a minimum of 10 gallons of spray suspension per acre by ground sprayer or in a minimum of 5 gallons of spray suspension per acre by aircraft.
				General directions: Observe fields closely for early disease symptoms. For optimum benefit, the lowest specified rate of a spray surfactant should be tank mixed with this product.
CROP	PHI (DAYS)	DISEASES	RATE OF PRODUCT PER ACRE	APPLICATION DIRECTIONS
Sunflowers	50	Rust (<i>Puccinia spp.</i>)	6 fl oz per acre Maximum 16 fl oz per crop season	Make a foliar spray application of this product (6 fl oz per acre) at the first sign of rust infection. Repeat this application after 14 days if the severity of the infection warrants treatment or if conditions conducive to infection prevail. Do not exceed the maximum specified treatment rate (16 fl oz) per crop season.
				Apply the specified treatment rate in a minimum quantity of 20 gallons of water per acre when making application of this product with ground sprayers. When making applications by aircraft use a minimum quantity of 5 gallons of water to disperse the specified application rate. Do not exceed the maximum specified application rate per crop per season.
General directions: Observe fields closely for early disease symptoms. For optimum benefit, the lowest specified rate of a spray surfactant should be tank mixed with this product. Thorough coverage is important for optimum disease control.				

NOTES: Follow all applicable directions, restrictions and precautions on the Muscle 3.6F product label.

This label must be in the possession of the user at the time of pesticide application on sunflowers or soybeans.

This supplemental label expires on June 30, 2011 and must not be used or distributed after this date.

Muscle is a registered trademark of ISAGRO S.p.A., Milan, Italy.

(100109) 1/10