

# Supplemental Labeling for Field Corn, Field Corn Grown for Seed, Popcorn

## RESTRICTED USE PESTICIDE

Toxic to fish and aquatic organisms

For retail sale to and use only by certified applicators, or persons under their direct supervision and only for those uses covered by the certified applicator's certification.



EPA Reg. No. 279-3426

**Active Ingredient:**

Zeta-Cypermethrin, \* S-Cyano (3-phenoxy-phenyl)methyl (+) cis/trans 3-(2,2-dichloroethenyl)-2,2 dimethylcyclopropane carboxylate.....9.15%

**Other Ingredients\*\*** ..... 90.85%  
100.0%

Contains 0.8 pounds active ingredient per gallon.

\* Cis/trans ratio: Max. 75% (±) cis and min. 25% (±) trans

\*\* Contains Petroleum Distillates

### KEEP OUT OF REACH OF CHILDREN WARNING AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

**It is a violation of Federal law to use this product in a manner inconsistent with its labeling.**

**This supplemental label expires on July 31, 2018 and must not be used or distributed after this date.**

**ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL ARE TO BE FOLLOWED.**

**This Supplemental labeling and the complete Mustang Maxx Insecticide label must be in possession of the user at the time of pesticide application. Read the label affixed to the container for Mustang Maxx Insecticide before applying. Carefully follow all precautionary statements and application use directions. Use of Mustang Maxx Insecticide according to this supplemental labeling is subject to all use precautions and limitations imposed by the labeling affixed to the container for Mustang Maxx Insecticide.**



Sold By:  
FMC Corporation  
Agricultural Products Group  
1735 Market Street  
Philadelphia, PA 19103

**Corn (Field), Field Corn Grown for Seed, Popcorn**

Insects Controlled	Rate of Application	Method of Application
Cutworms	1.28 to 2.8 fl oz/A (0.008 to 0.0175 lbs ai/A)	Make applications when insect populations reach economic thresholds. Refer to local Cooperative Extension Pest Management Guidelines and/or scouting results. Do not exceed maximum allowable rate.
Corn Earworm <sup>1</sup> Green Cloverworm Meadow Spittlebug Western Bean Cutworm <sup>1</sup>	1.76 to 4.0 fl oz/A (0.011 to 0.025 lbs ai/A)	
Bean Leaf Beetle Cereal Leaf Beetle Corn Borer, European Corn Borer, Southwestern Corn Rootworm Beetle Flea Beetle Grasshoppers Hop Vine Borer Hornworms Japanese Beetle (adult) Sap Beetle (adult) Southern Corn Leaf Beetle Stalk Borer Stink Bug spp. Tobacco Budworm <sup>2</sup> Webworms Aphids <sup>3</sup>	2.72 to 4.0 fl oz/A (0.017 to 0.025 lbs ai/A)	
Armyworms (including Fall Armyworms) Chinch Bug	3.2 to 4.0 fl oz/A (0.02 to 0.025 lbs ai/A)	
<p>Do not apply more than 16 fluid ounces of product (0.10 pound of active ingredient) per acre per season including at-plant plus foliar applications of Mustang Maxx Insecticide.</p> <p>Do not apply within 7 days of harvest for grain, stover and forage.</p> <p><sup>1</sup> For control before the larva bores into the plant stalk or ear.  <sup>2</sup> See resistance statement under "Directions for Use" section on main label.  <sup>3</sup> Control may be variable depending on species present and host-plant relationships.</p>		