



MANUFACTURED BY:
NACHURS ALPINE SOLUTIONS®
 421 Leader Street • Marion, OH 43302
 (800) 622-4877

visit us online:
www.nachurs.com

LOT #: _____

Bulk Shipment

See Bill of Lading for
 Volume & Net Weight

Jug: 2.5 U.S. Gallons (9.46 L)
 Net Weight: 24.75 LBS (11.23 kg)

Box: 2 x 2.5 U.S. Gallons
 (2 x 9.46 L)
 Net Weight: 49.5 LBS (22.45 kg)

Tote:
 Volume: _____
 Net Weight: _____

NACHURS® 3% Ca EDTA

0-0-7

**EDTA CHELATED CALCIUM FOR THE PREVENTION
 AND CORRECTION OF NUTRIENT DEFICIENCY**

GUARANTEED ANALYSIS

SOLUBLE POTASH (K₂O) 7%
 CALCIUM (Ca) 3%
 3% chelated calcium (Ca)

Derived from: potassium hydroxide and calcium EDTA.

DENSITY/ WEIGHT PER GALLON: 9.9 LBS (4.49 kg) at 68°F
 ALL COMPLETELY WATER SOLUBLE

PROTECT FROM FREEZING

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.htm>

USE IN ACCORDANCE WITH RECOMMENDATIONS OF A QUALIFIED INDIVIDUAL OR INSTITUTION, SUCH AS, BUT NOT LIMITED TO, A CERTIFIED CROP ADVISOR, AGRONOMIST, UNIVERSITY PUBLICATION OR APPLY ACCORDING TO RECOMMENDATIONS IN YOUR APPROVED NUTRIENT MANAGEMENT PLAN. NACHURS STARTER OR FOLIAR APPLICATIONS ARE INTENDED TO SUPPLEMENT EXISTING SOIL FERTILITY PROGRAMS AND WILL NOT BY ITSELF PROVIDE ALL THE NUTRIENTS NORMALLY REQUIRED BY AGRICULTURAL CROPS.

HAZARD STATEMENTS: HARMFUL TO AQUATIC LIFE. HARMFUL TO AQUATIC LIFE WITH LONG LASTING EFFECTS.

PRECAUTIONARY STATEMENTS: AVOID RELEASE TO THE ENVIRONMENT. DISPOSE OF CONTENTS/CONTAINER IN ACCORDANCE WITH LOCAL, REGIONAL, NATIONAL, AND INTERNATIONAL REGULATIONS.

WARNING: EXPOSURE MAY AGGRAVATE PRE-EXISTING EYE, SKIN, OR RESPIRATORY CONDITIONS.

"NACHURS ALPINE SOLUTIONS", "NACHURS", and "NACHURS" are trademarks of NACHURS ALPINE SOLUTIONS.

© 2016. NACHURS ALPINE SOLUTIONS.
 All rights reserved. FL- F000529



NACHURS® 3% Ca EDTA

EDTA CHELATED CALCIUM FOR THE PREVENTION AND CORRECTION OF NUTRIENT DEFICIENCY

MIXING INSTRUCTIONS

- Put 1/3 of water or fertilizer in tank
- Add other chemicals (if applicable)
- Fill tank with balance of water or fertilizer
- Add correct amount of chelate (shake each container vigorously before use)
- Agitate adequately to mix

CAUTION: Check compatibility with orthophosphates, high potash suspensions, and chemical mixtures.

GENERAL RECOMMENDATIONS:

NACHURS 3% Ca EDTA is intended for use in a complete crop nutritional program where a soil and/or plant tissue analysis has indicated a calcium deficiency. NACHURS 3% Ca EDTA may be applied with NACHURS liquid fertilizers, other liquid fertilizers, fertilizer suspensions, nitrogen solutions, or water.

| | pints/A | |
|---|------------------|---------------------|
| HIGH RESPONSE CROPS: | Row ¹ | Banded ² |
| alfalfa, potatoes, sugar beets, tobacco, tomatoes, clover, peppers, watermelon, lettuce, cabbage, brussel sprouts | 1-3 | 2-5 |
| OTHER CROPS: | 1-2 | 2-4 |
| corn, wheat, grasses, barley, oats, rye, sorghum | | |

Note: 1 pint of NACHURS 3% Ca EDTA contain: 0.037 pounds of elemental calcium.

- Row recommendations are intended to provide starter application of calcium for the current year crop and are not expected to correct severe calcium deficiencies. Additional banded or foliar applications may be necessary to correct severe deficiencies.
- Banded generally refers to 2" to the side and 2" below the seed and is usually applied during planting.

FOLIAR APPLICATIONS:

For all the above crops plus tree fruits, citrus, small fruits, and ornamentals:
 1-2 pints/A in minimum 10 gallons of water

Aerial: 1-2 pints/A in minimum 5 gallons of water

NACHURS liquid fertilizer may be used as a carrier at 3-5 gallons per acre. More than one application may be required for severe deficiency.

NOTE: Foliar fertilization is intended as a supplement to a regular fertilization program and will not by itself provide all the nutrients normally required by agricultural crops.

NACHURS chelated micronutrients should not be used unless a deficiency is indicated by a soil test and/or tissue analysis, and/or appearance of deficiency symptoms. Optimum rate of application will vary between fields depending on soil pH and organic matter content.



NACHURS® 3% Ca EDTA

EDTA CHELATED CALCIUM FOR THE PREVENTION AND CORRECTION OF NUTRIENT DEFICIENCY

MIXING INSTRUCTIONS

1. Put 1/3 of water or fertilizer in tank
2. Add other chemicals (if applicable)
3. Fill tank with balance of water or fertilizer
4. Add correct amount of chelate (shake each container vigorously before use)
5. Agitate adequately to mix

CAUTION: Check compatibility with orthophosphates, high potash suspensions, and chemical mixtures.

GENERAL RECOMMENDATIONS:

NACHURS 3% Ca EDTA is intended for use in a complete crop nutritional program where a soil and/or plant tissue analysis has indicated a calcium deficiency. NACHURS 3% Ca EDTA may be applied with NACHURS liquid fertilizers, other liquid fertilizers, fertilizer suspensions, nitrogen solutions, or water.

| | pints/A | |
|--|------------------|---------------------|
| | Row ¹ | Banded ² |
| HIGH RESPONSE CROPS: alfalfa, potatoes, sugar beets, tobacco, tomatoes, clover, peppers, watermelon, lettuce, cabbage, brussel sprouts | 1-3 | 2-5 |
| OTHER CROPS: corn, wheat, grasses, barley, oats, rye, sorghum | 1-2 | 2-4 |

Note: 1 pint of NACHURS 3% Ca EDTA contain: 0.037 pounds of elemental calcium.

1. Row recommendations are intended to provide starter application of calcium for the current year crop and are not expected to correct severe calcium deficiencies. Additional banded or foliar applications may be necessary to correct severe deficiencies.
2. Banded generally refers to 2" to the side and 2" below the seed and is usually applied during planting.

FOLIAR APPLICATIONS:

For all the above crops plus tree fruits, citrus, small fruits, and ornamentals:
1-2 pints/A in minimum 10 gallons of water

Aerial: 1-2 pints/A in minimum 5 gallons of water

NACHURS liquid fertilizer may be used as a carrier at 3-5 gallons per acre. More than one application may be required for severe deficiency.

NOTE: Foliar fertilization is intended as a supplement to a regular fertilization program and will not by itself provide all the nutrients normally required by agricultural crops.

NACHURS chelated micronutrients should not be used unless a deficiency is indicated by a soil test and/or tissue analysis, and/or appearance of deficiency symptoms. Optimum rate of application will vary between fields depending on soil pH and organic matter content.



NACHURS® 3% Ca EDTA

EDTA CHELATED CALCIUM FOR THE PREVENTION AND CORRECTION OF NUTRIENT DEFICIENCY

MIXING INSTRUCTIONS

1. Put 1/3 of water or fertilizer in tank
2. Add other chemicals (if applicable)
3. Fill tank with balance of water or fertilizer
4. Add correct amount of chelate (shake each container vigorously before use)
5. Agitate adequately to mix

CAUTION: Check compatibility with orthophosphates, high potash suspensions, and chemical mixtures.

GENERAL RECOMMENDATIONS:

NACHURS 3% Ca EDTA is intended for use in a complete crop nutritional program where a soil and/or plant tissue analysis has indicated a calcium deficiency. NACHURS 3% Ca EDTA may be applied with NACHURS liquid fertilizers, other liquid fertilizers, fertilizer suspensions, nitrogen solutions, or water.

| | pints/A | |
|--|------------------|---------------------|
| | Row ¹ | Banded ² |
| HIGH RESPONSE CROPS: alfalfa, potatoes, sugar beets, tobacco, tomatoes, clover, peppers, watermelon, lettuce, cabbage, brussel sprouts | 1-3 | 2-5 |
| OTHER CROPS: corn, wheat, grasses, barley, oats, rye, sorghum | 1-2 | 2-4 |

Note: 1 pint of NACHURS 3% Ca EDTA contain: 0.037 pounds of elemental calcium.

1. Row recommendations are intended to provide starter application of calcium for the current year crop and are not expected to correct severe calcium deficiencies. Additional banded or foliar applications may be necessary to correct severe deficiencies.
2. Banded generally refers to 2" to the side and 2" below the seed and is usually applied during planting.

FOLIAR APPLICATIONS:

For all the above crops plus tree fruits, citrus, small fruits, and ornamentals:
1-2 pints/A in minimum 10 gallons of water

Aerial: 1-2 pints/A in minimum 5 gallons of water

NACHURS liquid fertilizer may be used as a carrier at 3-5 gallons per acre. More than one application may be required for severe deficiency.

NOTE: Foliar fertilization is intended as a supplement to a regular fertilization program and will not by itself provide all the nutrients normally required by agricultural crops.

NACHURS chelated micronutrients should not be used unless a deficiency is indicated by a soil test and/or tissue analysis, and/or appearance of deficiency symptoms. Optimum rate of application will vary between fields depending on soil pH and organic matter content.