



MANUFACTURED BY:  
**NACHURS ALPINE SOLUTIONS®**  
 421 Leader Street • Marion, OH 43302  
 (800) 622-4877

visit us online:  
[www.nachurs.com](http://www.nachurs.com)

LOT #: \_\_\_\_\_

**Bulk Shipment**

See Bill of Lading for  
 Volume & Net Weight

**Jug:** 2.5 U.S. Gallons (9.46 L)  
 Net Weight: 26.28 LBS (11.92 kg)

**Box:** 2 x 2.5 U.S. Gallons  
 (2 x 9.46 L)  
 Net Weight: 52.55 LBS (23.84 kg)

**Tote:**  
 Volume: \_\_\_\_\_  
 Net Weight: \_\_\_\_\_

**NACHURS® 7.5% Cu EDTA**

**3-0-0**

**EDTA CHELATED COPPER FOR THE PREVENTION  
 AND CORRECTION OF NUTRIENT DEFICIENCY**

**GUARANTEED ANALYSIS**

TOTAL NITROGEN (N) ..... 3%  
 3% ammoniacal nitrogen  
 COPPER (Cu) ..... 7.5%  
 7.5% chelated copper (Cu)

Derived from: ammonium hydroxide and copper EDTA.

**DENSITY/ WEIGHT PER GALLON: 10.51 LBS (4.77 kg) at 68°F**  
 ALL COMPLETELY WATER SOLUBLE

PROTECT FROM FREEZING

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.htm>

USE IN ACCORDANCE WITH RECOMMENDATIONS OF A QUALIFIED INDIVIDUAL OR INSTITUTION, SUCH AS, BUT NOT LIMITED TO, A CERTIFIED CROP ADVISOR, AGRONOMIST, UNIVERSITY PUBLICATION OR APPLY ACCORDING TO RECOMMENDATIONS IN YOUR APPROVED NUTRIENT MANAGEMENT PLAN. NACHURS STARTER OR FOLIAR APPLICATIONS ARE INTENDED TO SUPPLEMENT EXISTING SOIL FERTILITY PROGRAMS AND WILL NOT BY ITSELF PROVIDE ALL THE NUTRIENTS NORMALLY REQUIRED BY AGRICULTURAL CROPS.



WARNING

**HAZARD STATEMENTS:** HARMFUL IF SWALLOWED. CAUSES SKIN IRRITATION. CAUSES SERIOUS EYE IRRITATION.  
**PRECAUTIONARY STATEMENTS:** WASH HANDS, FOREARMS, AND OTHER EXPOSED AREAS THOROUGHLY AFTER HANDLING. DO NOT EAT, DRINK OR SMOKE WHEN USING THIS PRODUCT. WEAR PROTECTIVE GLOVES, PROTECTIVE CLOTHING, AND EYE PROTECTION. IF SWALLOWED: CALL A POISON CENTER OR DOCTOR IF YOU FEEL UNWELL. IF ON SKIN: WASH WITH PLENTY OF WATER. IF IN EYES: RINSE CAUTIOUSLY WITH WATER FOR SEVERAL MINUTES. REMOVE CONTACT LENSES, IF PRESENT AND EASY TO DO. CONTINUE RINSING. SPECIFIC TREATMENT-SEE SECTION 4 OF THE SDS. RINSE MOUTH. IF SKIN IRRITATION OCCURS: GET MEDICAL ADVICE/ATTENTION. IF EYE IRRITATION PERSISTS: GET MEDICAL ADVICE/ATTENTION. TAKE OFF CONTAMINATED CLOTHING AND WASH IT BEFORE REUSE. DISPOSE OF CONTENTS/CONTAINER IN ACCORDANCE WITH LOCAL, REGIONAL, NATIONAL, AND INTERNATIONAL REGULATIONS.  
**WARNING:** EXPOSURE MAY AGGRAVATE PRE-EXISTING EYE, SKIN, OR RESPIRATORY CONDITIONS.

"NACHURS ALPINE SOLUTIONS", "aces", and "NACHURS" are trademarks of NACHURS ALPINE SOLUTIONS.

© 2016. NACHURS ALPINE SOLUTIONS.  
 All rights reserved. FL- F000529



**NACHURS® 7.5% Cu EDTA**

**EDTA CHELATED COPPER FOR THE PREVENTION AND CORRECTION OF NUTRIENT DEFICIENCY**

**MIXING INSTRUCTIONS**

- Put 1/3 of water or fertilizer in tank
- Add other chemicals (if applicable)
- Fill tank with balance of water or fertilizer
- Add correct amount of chelate (shake each container vigorously before use)
- Agitate adequately to mix

**CAUTION:** Check compatibility with orthophosphates, high potash suspensions, and chemical mixtures.

**GENERAL RECOMMENDATIONS:**

NACHURS 7.5% Cu EDTA is intended for use in a complete crop nutritional program where a soil and/or plant tissue analysis has indicated a copper deficiency. NACHURS 7.5% Cu EDTA may be applied with NACHURS liquid fertilizers, other liquid fertilizers, fertilizer suspensions, nitrogen solutions, or water.

	pints/A	
HIGH RESPONSE CROPS:	Row <sup>1</sup>	Banded <sup>2</sup>
alfalfa, potatoes, sugar beets, tobacco, tomatoes, clover, peppers, watermelon, lettuce, cabbage, brussel sprouts	1-3	2-5
OTHER CROPS:	1-2	2-4
corn, wheat, grasses, barley, oats, rye, sorghum		

Note: 1 pint of NACHURS 7.5% Cu EDTA contains: 0.099 pounds of elemental copper.

- Row recommendations are intended to provide starter application of copper for the current year crop and are not expected to correct severe copper deficiencies. Additional banded or foliar applications may be necessary to correct severe deficiencies.
- Banded generally refers to 2" to the side and 2" below the seed and is usually applied during planting.

**FOLIAR APPLICATIONS:**

**For all the above crops plus tree fruits, citrus, small fruits, and ornamentals:**  
 1-2 pints/A in minimum 10 gallons of water

**Aerial:** 1-2 pints/A in minimum 5 gallons of water

NACHURS liquid fertilizer may be used as a carrier at 3-5 gallons per acre. More than one application may be required for severe deficiency.

**NOTE:** Foliar fertilization is intended as a supplement to a regular fertilization program and will not by itself provide all the nutrients normally required by agricultural crops.

**NACHURS chelated micronutrients should not be used unless a deficiency is indicated by a soil test and/or tissue analysis, and/or appearance of deficiency symptoms. Optimum rate of application will vary between fields depending on soil pH and organic matter content.**



MANUFACTURED BY:  
**NACHURS ALPINE SOLUTIONS®**  
 421 Leader Street • Marion, OH 43302  
 (800) 622-4877

visit us online:  
[www.nachurs.com](http://www.nachurs.com)

LOT #: \_\_\_\_\_

**Bulk Shipment**

See Bill of Lading for  
 Volume & Net Weight

**Jug:** 2.5 U.S. Gallons (9.46 L)  
 Net Weight: 26.28 LBS (11.92 kg)

**Box:** 2 x 2.5 U.S. Gallons  
 (2 x 9.46 L)  
 Net Weight: 52.55 LBS (23.84 kg)

**Tote:**  
 Volume: \_\_\_\_\_  
 Net Weight: \_\_\_\_\_

**NACHURS® 7.5% Cu EDTA**

**3-0-0**

**EDTA CHELATED COPPER FOR THE PREVENTION  
 AND CORRECTION OF NUTRIENT DEFICIENCY**

**GUARANTEED ANALYSIS**

TOTAL NITROGEN (N) .....	3%
3% ammoniacal nitrogen	
COPPER (Cu) .....	7.5%
7.5% chelated copper (Cu)	

Derived from: ammonium hydroxide and copper EDTA.

**DENSITY/ WEIGHT PER GALLON: 10.51 LBS (4.77 kg) at 68°F**  
 ALL COMPLETELY WATER SOLUBLE

PROTECT FROM FREEZING

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfc.org/metals.htm>

USE IN ACCORDANCE WITH RECOMMENDATIONS OF A QUALIFIED INDIVIDUAL OR INSTITUTION, SUCH AS, BUT NOT LIMITED TO, A CERTIFIED CROP ADVISOR, AGRONOMIST, UNIVERSITY PUBLICATION OR APPLY ACCORDING TO RECOMMENDATIONS IN YOUR APPROVED NUTRIENT MANAGEMENT PLAN. NACHURS STARTER OR FOLIAR APPLICATIONS ARE INTENDED TO SUPPLEMENT EXISTING SOIL FERTILITY PROGRAMS AND WILL NOT BY ITSELF PROVIDE ALL THE NUTRIENTS NORMALLY REQUIRED BY AGRICULTURAL CROPS.



WARNING

**HAZARD STATEMENTS:** HARMFUL IF SWALLOWED. CAUSES SKIN IRRITATION. CAUSES SERIOUS EYE IRRITATION.  
**PRECAUTIONARY STATEMENTS:** WASH HANDS, FOREARMS, AND OTHER EXPOSED AREAS THOROUGHLY AFTER HANDLING. DO NOT EAT, DRINK OR SMOKE WHEN USING THIS PRODUCT. WEAR PROTECTIVE GLOVES, PROTECTIVE CLOTHING, AND EYE PROTECTION. IF SWALLOWED: CALL A POISON CENTER OR DOCTOR IF YOU FEEL UNWELL. IF ON SKIN: WASH WITH PLENTY OF WATER. IF IN EYES: RINSE CAUTIOUSLY WITH WATER FOR SEVERAL MINUTES. REMOVE CONTACT LENSES, IF PRESENT AND EASY TO DO. CONTINUE RINSING. SPECIFIC TREATMENT-SEE SECTION 4 OF THE SDS. RINSE MOUTH. IF SKIN IRRITATION OCCURS: GET MEDICAL ADVICE/ATTENTION. IF EYE IRRITATION PERSISTS: GET MEDICAL ADVICE/ATTENTION. TAKE OFF CONTAMINATED CLOTHING AND WASH IT BEFORE REUSE. DISPOSE OF CONTENTS/CONTAINER IN ACCORDANCE WITH LOCAL, REGIONAL, NATIONAL, AND INTERNATIONAL REGULATIONS.  
**WARNING:** EXPOSURE MAY AGGRAVATE PRE-EXISTING EYE, SKIN, OR RESPIRATORY CONDITIONS.

"NACHURS ALPINE SOLUTIONS", "aces", and "NACHURS" are trademarks of NACHURS ALPINE SOLUTIONS.

© 2016. NACHURS ALPINE SOLUTIONS.  
 All rights reserved. FL- F000529



MANUFACTURED BY:  
**NACHURS ALPINE SOLUTIONS®**  
 421 Leader Street • Marion, OH 43302  
 (800) 622-4877

visit us online:  
[www.nachurs.com](http://www.nachurs.com)

LOT #: \_\_\_\_\_

**Bulk Shipment**

See Bill of Lading for  
 Volume & Net Weight

**Jug:** 2.5 U.S. Gallons (9.46 L)  
 Net Weight: 26.28 LBS (11.92 kg)

**Box:** 2 x 2.5 U.S. Gallons  
 (2 x 9.46 L)  
 Net Weight: 52.55 LBS (23.84 kg)

**Tote:**  
 Volume: \_\_\_\_\_  
 Net Weight: \_\_\_\_\_

**NACHURS® 7.5% Cu EDTA**

**3-0-0**

**EDTA CHELATED COPPER FOR THE PREVENTION  
 AND CORRECTION OF NUTRIENT DEFICIENCY**

**GUARANTEED ANALYSIS**

TOTAL NITROGEN (N) .....	3%
3% ammoniacal nitrogen	
COPPER (Cu) .....	7.5%
7.5% chelated copper (Cu)	

Derived from: ammonium hydroxide and copper EDTA.

**DENSITY/ WEIGHT PER GALLON: 10.51 LBS (4.77 kg) at 68°F**  
 ALL COMPLETELY WATER SOLUBLE

PROTECT FROM FREEZING

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfc.org/metals.htm>

USE IN ACCORDANCE WITH RECOMMENDATIONS OF A QUALIFIED INDIVIDUAL OR INSTITUTION, SUCH AS, BUT NOT LIMITED TO, A CERTIFIED CROP ADVISOR, AGRONOMIST, UNIVERSITY PUBLICATION OR APPLY ACCORDING TO RECOMMENDATIONS IN YOUR APPROVED NUTRIENT MANAGEMENT PLAN. NACHURS STARTER OR FOLIAR APPLICATIONS ARE INTENDED TO SUPPLEMENT EXISTING SOIL FERTILITY PROGRAMS AND WILL NOT BY ITSELF PROVIDE ALL THE NUTRIENTS NORMALLY REQUIRED BY AGRICULTURAL CROPS.



WARNING

**HAZARD STATEMENTS:** HARMFUL IF SWALLOWED. CAUSES SKIN IRRITATION. CAUSES SERIOUS EYE IRRITATION.  
**PRECAUTIONARY STATEMENTS:** WASH HANDS, FOREARMS, AND OTHER EXPOSED AREAS THOROUGHLY AFTER HANDLING. DO NOT EAT, DRINK OR SMOKE WHEN USING THIS PRODUCT. WEAR PROTECTIVE GLOVES, PROTECTIVE CLOTHING, AND EYE PROTECTION. IF SWALLOWED: CALL A POISON CENTER OR DOCTOR IF YOU FEEL UNWELL. IF ON SKIN: WASH WITH PLENTY OF WATER. IF IN EYES: RINSE CAUTIOUSLY WITH WATER FOR SEVERAL MINUTES. REMOVE CONTACT LENSES, IF PRESENT AND EASY TO DO. CONTINUE RINSING. SPECIFIC TREATMENT-SEE SECTION 4 OF THE SDS. RINSE MOUTH. IF SKIN IRRITATION OCCURS: GET MEDICAL ADVICE/ATTENTION. IF EYE IRRITATION PERSISTS: GET MEDICAL ADVICE/ATTENTION. TAKE OFF CONTAMINATED CLOTHING AND WASH IT BEFORE REUSE. DISPOSE OF CONTENTS/CONTAINER IN ACCORDANCE WITH LOCAL, REGIONAL, NATIONAL, AND INTERNATIONAL REGULATIONS.  
**WARNING:** EXPOSURE MAY AGGRAVATE PRE-EXISTING EYE, SKIN, OR RESPIRATORY CONDITIONS.

"NACHURS ALPINE SOLUTIONS", "aces", and "NACHURS" are trademarks of NACHURS ALPINE SOLUTIONS.

© 2016. NACHURS ALPINE SOLUTIONS.  
 All rights reserved. FL- F000529



# NACHURS® 7.5% Cu EDTA

EDTA CHELATED COPPER FOR THE PREVENTION AND CORRECTION OF NUTRIENT DEFICIENCY

### MIXING INSTRUCTIONS

1. Put 1/3 of water or fertilizer in tank
2. Add other chemicals (if applicable)
3. Fill tank with balance of water or fertilizer
4. Add correct amount of chelate (shake each container vigorously before use)
5. Agitate adequately to mix

**CAUTION:** Check compatibility with orthophosphates, high potash suspensions, and chemical mixtures.

### GENERAL RECOMMENDATIONS:

NACHURS 7.5% Cu EDTA is intended for use in a complete crop nutritional program where a soil and/or plant tissue analysis has indicated a copper deficiency. NACHURS 7.5% Cu EDTA may be applied with NACHURS liquid fertilizers, other liquid fertilizers, fertilizer suspensions, nitrogen solutions, or water.

	pints/A	
	Row <sup>1</sup>	Banded <sup>2</sup>
<b>HIGH RESPONSE CROPS:</b> alfalfa, potatoes, sugar beets, tobacco, tomatoes, clover, peppers, watermelon, lettuce, cabbage, brussel sprouts	1-3	2-5
<b>OTHER CROPS:</b> corn, wheat, grasses, barley, oats, rye, sorghum	1-2	2-4

Note: 1 pint of NACHURS 7.5% Cu EDTA contains: 0.099 pounds of elemental copper.

1. Row recommendations are intended to provide starter application of copper for the current year crop and are not expected to correct severe copper deficiencies. Additional banded or foliar applications may be necessary to correct severe deficiencies.
2. Banded generally refers to 2" to the side and 2" below the seed and is usually applied during planting.

### FOLIAR APPLICATIONS:

**For all the above crops plus tree fruits, citrus, small fruits, and ornamentals:**  
1-2 pints/A in minimum 10 gallons of water

**Aerial:** 1-2 pints/A in minimum 5 gallons of water

NACHURS liquid fertilizer may be used as a carrier at 3-5 gallons per acre. More than one application may be required for severe deficiency.

**NOTE:** Foliar fertilization is intended as a supplement to a regular fertilization program and will not by itself provide all the nutrients normally required by agricultural crops.

**NACHURS chelated micronutrients should not be used unless a deficiency is indicated by a soil test and/or tissue analysis, and/or appearance of deficiency symptoms. Optimum rate of application will vary between fields depending on soil pH and organic matter content.**



# NACHURS® 7.5% Cu EDTA

EDTA CHELATED COPPER FOR THE PREVENTION AND CORRECTION OF NUTRIENT DEFICIENCY

### MIXING INSTRUCTIONS

1. Put 1/3 of water or fertilizer in tank
2. Add other chemicals (if applicable)
3. Fill tank with balance of water or fertilizer
4. Add correct amount of chelate (shake each container vigorously before use)
5. Agitate adequately to mix

**CAUTION:** Check compatibility with orthophosphates, high potash suspensions, and chemical mixtures.

### GENERAL RECOMMENDATIONS:

NACHURS 7.5% Cu EDTA is intended for use in a complete crop nutritional program where a soil and/or plant tissue analysis has indicated a copper deficiency. NACHURS 7.5% Cu EDTA may be applied with NACHURS liquid fertilizers, other liquid fertilizers, fertilizer suspensions, nitrogen solutions, or water.

	pints/A	
	Row <sup>1</sup>	Banded <sup>2</sup>
<b>HIGH RESPONSE CROPS:</b> alfalfa, potatoes, sugar beets, tobacco, tomatoes, clover, peppers, watermelon, lettuce, cabbage, brussel sprouts	1-3	2-5
<b>OTHER CROPS:</b> corn, wheat, grasses, barley, oats, rye, sorghum	1-2	2-4

Note: 1 pint of NACHURS 7.5% Cu EDTA contains: 0.099 pounds of elemental copper.

1. Row recommendations are intended to provide starter application of copper for the current year crop and are not expected to correct severe copper deficiencies. Additional banded or foliar applications may be necessary to correct severe deficiencies.
2. Banded generally refers to 2" to the side and 2" below the seed and is usually applied during planting.

### FOLIAR APPLICATIONS:

**For all the above crops plus tree fruits, citrus, small fruits, and ornamentals:**  
1-2 pints/A in minimum 10 gallons of water

**Aerial:** 1-2 pints/A in minimum 5 gallons of water

NACHURS liquid fertilizer may be used as a carrier at 3-5 gallons per acre. More than one application may be required for severe deficiency.

**NOTE:** Foliar fertilization is intended as a supplement to a regular fertilization program and will not by itself provide all the nutrients normally required by agricultural crops.

**NACHURS chelated micronutrients should not be used unless a deficiency is indicated by a soil test and/or tissue analysis, and/or appearance of deficiency symptoms. Optimum rate of application will vary between fields depending on soil pH and organic matter content.**