



MANUFACTURED BY:
NACHURS ALPINE SOLUTIONS®
 421 Leader Street • Marion, OH 43302
 (800) 622-4877

visit us online:
www.nachurs.com

LOT #: _____

Bulk Shipment

See Bill of Lading for
 Volume & Net Weight

Jug: 2.5 U.S. Gallons (9.46 L)
 Net Weight: 25.55 LBS (11.59 kg)

Box: 2 x 2.5 U.S. Gallons
 (2 x 9.46 L)
 Net Weight: 51.1 LBS (23.18 kg)

Tote:
 Volume: _____
 Net Weight: _____

NACHURS CropMax®

2-0-2

**CHELATED MICRONUTRIENT PRE-MIX FOR
 PREVENTION AND CORRECTION OF MICRONUTRIENT DEFICIENCY IN ALL CROPS**

GUARANTEED ANALYSIS

TOTAL NITROGEN (N)	2%
2% Ammoniacal Nitrogen	
SOLUBLE POTASH (K ₂ O)	2%
BORON (B)	0.1%
COPPER (Cu)	0.15%
0.15% Chelated Copper (Cu)	
IRON (Fe)	0.3%
0.3% Chelated Iron (Fe)	
MANGANESE (Mn)	1.5%
1.5% Chelated Manganese (Mn)	
MOLYBDENUM (Mo)	0.0005%
ZINC (Zn)	4.0%
4.0% Chelated Zinc (Zn)	

Derived from: ammonium hydroxide, potassium hydroxide, boric acid, copper EDTA, iron EDTA, manganese EDTA, sodium molybdate, zinc EDTA

DENSITY/ WEIGHT PER GALLON: 10.22 LBS (4.64 kg) at 68°F
 ALL COMPLETELY WATER SOLUBLE

**THIS PRODUCT IS INTENDED FOR USE IN MIXING OR BLENDING WITH OTHER FERTILIZER MATERIALS,
 TO PRODUCE FERTILIZERS WHOSE TOTAL PRIMARY NUTRIENT GUARANTEES EQUAL OR EXCEED 24%**

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.htm>

WARNING: BORON IS TO BE USED WHERE SOIL TEST AND/OR TISSUE ANALYSIS INDICATE A DEFICIENCY AND SHOULD NOT BE USED AT RATES IN EXCESS OF THE RATE RECOMMENDED BY A QUALIFIED INDIVIDUAL/ENTITY SUCH AS A CERTIFIED CROP ADVISOR, AGRONOMIST OR UNIVERSITY. EXCESSIVE APPLICATION OF BORON MAY CAUSE CROP DAMAGE. THIS FERTILIZER IS TO BE USED ONLY ON SOIL WHICH RESPONDS TO MOLYBDENUM. CROPS HIGH IN MOLYBDENUM ARE TOXIC TO GRAZING ANIMALS (RUMINANTS).

USE IN ACCORDANCE WITH RECOMMENDATIONS OF A QUALIFIED INDIVIDUAL OR INSTITUTION, SUCH AS, BUT NOT LIMITED TO, A CERTIFIED CROP ADVISOR, AGRONOMIST, UNIVERSITY PUBLICATION OR APPLY ACCORDING TO RECOMMENDATIONS IN YOUR APPROVED NUTRIENT MANAGEMENT PLAN. NACHURS STARTER OR FOLIAR APPLICATIONS ARE INTENDED TO SUPPLEMENT EXISTING SOIL FERTILITY PROGRAMS AND WILL NOT BY ITSELF PROVIDE ALL THE NUTRIENTS NORMALLY REQUIRED BY AGRICULTURAL CROPS.

WARNING: EXPOSURE MAY AGGRAVATE PRE-EXISTING EYE, SKIN, OR RESPIRATORY CONDITIONS.

"NACHURS ALPINE SOLUTIONS", "NACHURS", and "NACHURS CropMax" are trademarks of NACHURS ALPINE SOLUTIONS.

© 2018. NACHURS ALPINE SOLUTIONS.
 All rights reserved. FL- F000529



NACHURS CropMax®

CHELATED MICRONUTRIENT PRE-MIX FOR THE PREVENTION AND CORRECTION OF MICRONUTRIENT DEFICIENCY

APPLICATION INSTRUCTIONS:

Suggested application rate: 1-2 pts/acre.

NACHURS CropMax is formulated to provide necessary micronutrients for optimum growth with the advantage of excellent compatibility with NACHURS in-furrow starters.

Compatibility: **NACHURS CropMax** may be applied in combination with other liquid fertilizers, fertilizer suspensions, and nitrogen solutions. Always jar test combinations before field mixing.

GENERAL MIXING INSTRUCTIONS:

- 1) Put 1/3 of fertilizer in tank
- 2) Add other chemicals, if any
- 3) Fill tank with balance of fertilizer
- 4) Add correct amount of chelate
- 5) Agitate adequately to mix

CAUTION: Check compatibility with standard jar test.

NACHURS CropMax must be applied with NACHURS liquid fertilizers, other liquid fertilizers, fertilizer suspensions, and nitrogen solutions.

These are general product recommendations. Please consult with your NACHURS Sales Manager or agronomist for specific fertility recommendations.



MANUFACTURED BY:
NACHURS ALPINE SOLUTIONS®
 421 Leader Street • Marion, OH 43302
 (800) 622-4877

visit us online:
www.nachurs.com

LOT #: _____

Bulk Shipment

See Bill of Lading for
 Volume & Net Weight

Jug: 2.5 U.S. Gallons (9.46 L)
 Net Weight: 25.55 LBS (11.59 kg)

Box: 2 x 2.5 U.S. Gallons
 (2 x 9.46 L)
 Net Weight: 51.1 LBS (23.18 kg)

Tote:
 Volume: _____
 Net Weight: _____

NACHURS CropMax®

2-0-2

**CHELATED MICRONUTRIENT PRE-MIX FOR
 PREVENTION AND CORRECTION OF MICRONUTRIENT DEFICIENCY IN ALL CROPS**

GUARANTEED ANALYSIS

TOTAL NITROGEN (N)	2%
2% Ammoniacal Nitrogen	
SOLUBLE POTASH (K ₂ O)	2%
BORON (B)	0.1%
COPPER (Cu)	0.15%
0.15% Chelated Copper (Cu)	
IRON (Fe)	0.3%
0.3% Chelated Iron (Fe)	
MANGANESE (Mn)	1.5%
1.5% Chelated Manganese (Mn)	
MOLYBDENUM (Mo)	0.0005%
ZINC (Zn)	4.0%
4.0% Chelated Zinc (Zn)	

Derived from: ammonium hydroxide, potassium hydroxide, boric acid, copper EDTA, iron EDTA, manganese EDTA, sodium molybdate, zinc EDTA

DENSITY/ WEIGHT PER GALLON: 10.22 LBS (4.64 kg) at 68°F
 ALL COMPLETELY WATER SOLUBLE

**THIS PRODUCT IS INTENDED FOR USE IN MIXING OR BLENDING WITH OTHER FERTILIZER MATERIALS,
 TO PRODUCE FERTILIZERS WHOSE TOTAL PRIMARY NUTRIENT GUARANTEES EQUAL OR EXCEED 24%**

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.htm>

WARNING: BORON IS TO BE USED WHERE SOIL TEST AND/OR TISSUE ANALYSIS INDICATE A DEFICIENCY AND SHOULD NOT BE USED AT RATES IN EXCESS OF THE RATE RECOMMENDED BY A QUALIFIED INDIVIDUAL/ENTITY SUCH AS A CERTIFIED CROP ADVISOR, AGRONOMIST OR UNIVERSITY. EXCESSIVE APPLICATION OF BORON MAY CAUSE CROP DAMAGE. THIS FERTILIZER IS TO BE USED ONLY ON SOIL WHICH RESPONDS TO MOLYBDENUM. CROPS HIGH IN MOLYBDENUM ARE TOXIC TO GRAZING ANIMALS (RUMINANTS).

USE IN ACCORDANCE WITH RECOMMENDATIONS OF A QUALIFIED INDIVIDUAL OR INSTITUTION, SUCH AS, BUT NOT LIMITED TO, A CERTIFIED CROP ADVISOR, AGRONOMIST, UNIVERSITY PUBLICATION OR APPLY ACCORDING TO RECOMMENDATIONS IN YOUR APPROVED NUTRIENT MANAGEMENT PLAN. NACHURS STARTER OR FOLIAR APPLICATIONS ARE INTENDED TO SUPPLEMENT EXISTING SOIL FERTILITY PROGRAMS AND WILL NOT BY ITSELF PROVIDE ALL THE NUTRIENTS NORMALLY REQUIRED BY AGRICULTURAL CROPS.

WARNING: EXPOSURE MAY AGGRAVATE PRE-EXISTING EYE, SKIN, OR RESPIRATORY CONDITIONS.

"NACHURS ALPINE SOLUTIONS", "NACHURS", "NACHURS", and "NACHURS CropMax" are trademarks of NACHURS ALPINE SOLUTIONS.

© 2017. NACHURS ALPINE SOLUTIONS.
 All rights reserved. FL- F000529



MANUFACTURED BY:
NACHURS ALPINE SOLUTIONS®
 421 Leader Street • Marion, OH 43302
 (800) 622-4877

visit us online:
www.nachurs.com

LOT #: _____

Bulk Shipment

See Bill of Lading for
 Volume & Net Weight

Jug: 2.5 U.S. Gallons (9.46 L)
 Net Weight: 25.55 LBS (11.59 kg)

Box: 2 x 2.5 U.S. Gallons
 (2 x 9.46 L)
 Net Weight: 51.1 LBS (23.18 kg)

Tote:
 Volume: _____
 Net Weight: _____

NACHURS CropMax®

2-0-2

**CHELATED MICRONUTRIENT PRE-MIX FOR
 PREVENTION AND CORRECTION OF MICRONUTRIENT DEFICIENCY IN ALL CROPS**

GUARANTEED ANALYSIS

TOTAL NITROGEN (N)	2%
2% Ammoniacal Nitrogen	
SOLUBLE POTASH (K ₂ O)	2%
BORON (B)	0.1%
COPPER (Cu)	0.15%
0.15% Chelated Copper (Cu)	
IRON (Fe)	0.3%
0.3% Chelated Iron (Fe)	
MANGANESE (Mn)	1.5%
1.5% Chelated Manganese (Mn)	
MOLYBDENUM (Mo)	0.0005%
ZINC (Zn)	4.0%
4.0% Chelated Zinc (Zn)	

Derived from: ammonium hydroxide, potassium hydroxide, boric acid, copper EDTA, iron EDTA, manganese EDTA, sodium molybdate, zinc EDTA

DENSITY/ WEIGHT PER GALLON: 10.22 LBS (4.64 kg) at 68°F
 ALL COMPLETELY WATER SOLUBLE

**THIS PRODUCT IS INTENDED FOR USE IN MIXING OR BLENDING WITH OTHER FERTILIZER MATERIALS,
 TO PRODUCE FERTILIZERS WHOSE TOTAL PRIMARY NUTRIENT GUARANTEES EQUAL OR EXCEED 24%**

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.htm>

WARNING: BORON IS TO BE USED WHERE SOIL TEST AND/OR TISSUE ANALYSIS INDICATE A DEFICIENCY AND SHOULD NOT BE USED AT RATES IN EXCESS OF THE RATE RECOMMENDED BY A QUALIFIED INDIVIDUAL/ENTITY SUCH AS A CERTIFIED CROP ADVISOR, AGRONOMIST OR UNIVERSITY. EXCESSIVE APPLICATION OF BORON MAY CAUSE CROP DAMAGE. THIS FERTILIZER IS TO BE USED ONLY ON SOIL WHICH RESPONDS TO MOLYBDENUM. CROPS HIGH IN MOLYBDENUM ARE TOXIC TO GRAZING ANIMALS (RUMINANTS).

USE IN ACCORDANCE WITH RECOMMENDATIONS OF A QUALIFIED INDIVIDUAL OR INSTITUTION, SUCH AS, BUT NOT LIMITED TO, A CERTIFIED CROP ADVISOR, AGRONOMIST, UNIVERSITY PUBLICATION OR APPLY ACCORDING TO RECOMMENDATIONS IN YOUR APPROVED NUTRIENT MANAGEMENT PLAN. NACHURS STARTER OR FOLIAR APPLICATIONS ARE INTENDED TO SUPPLEMENT EXISTING SOIL FERTILITY PROGRAMS AND WILL NOT BY ITSELF PROVIDE ALL THE NUTRIENTS NORMALLY REQUIRED BY AGRICULTURAL CROPS.

WARNING: EXPOSURE MAY AGGRAVATE PRE-EXISTING EYE, SKIN, OR RESPIRATORY CONDITIONS.

"NACHURS ALPINE SOLUTIONS", "NACHURS", "NACHURS", and "NACHURS CropMax" are trademarks of NACHURS ALPINE SOLUTIONS.

© 2017. NACHURS ALPINE SOLUTIONS.
 All rights reserved. FL- F000529



NACHURS CropMax®

CHELATED MICRONUTRIENT PRE-MIX FOR THE PREVENTION AND CORRECTION OF MICRONUTRIENT DEFICIENCY

APPLICATION INSTRUCTIONS:

Suggested application rate: 1-2 pts/acre.

NACHURS CropMax is formulated to provide necessary micronutrients for optimum growth with the advantage of excellent compatibility with NACHURS in-furrow starters.

Compatibility: **NACHURS CropMax** may be applied in combination with other liquid fertilizers, fertilizer suspensions, and nitrogen solutions. Always jar test combinations before field mixing.

GENERAL MIXING INSTRUCTIONS:

- 1) Put 1/3 of fertilizer in tank
- 2) Add other chemicals, if any
- 3) Fill tank with balance of fertilizer
- 4) Add correct amount of chelate
- 5) Agitate adequately to mix

CAUTION: Check compatibility with standard jar test.

NACHURS CropMax must be applied with NACHURS liquid fertilizers, other liquid fertilizers, fertilizer suspensions, and nitrogen solutions.

These are general product recommendations. Please consult with your NACHURS Sales Manager or agronomist for specific fertility recommendations.



NACHURS CropMax®

CHELATED MICRONUTRIENT PRE-MIX FOR THE PREVENTION AND CORRECTION OF MICRONUTRIENT DEFICIENCY

APPLICATION INSTRUCTIONS:

Suggested application rate: 1-2 pts/acre.

NACHURS CropMax is formulated to provide necessary micronutrients for optimum growth with the advantage of excellent compatibility with NACHURS in-furrow starters.

Compatibility: **NACHURS CropMax** may be applied in combination with other liquid fertilizers, fertilizer suspensions, and nitrogen solutions. Always jar test combinations before field mixing.

GENERAL MIXING INSTRUCTIONS:

- 1) Put 1/3 of fertilizer in tank
- 2) Add other chemicals, if any
- 3) Fill tank with balance of fertilizer
- 4) Add correct amount of chelate
- 5) Agitate adequately to mix

CAUTION: Check compatibility with standard jar test.

NACHURS CropMax must be applied with NACHURS liquid fertilizers, other liquid fertilizers, fertilizer suspensions, and nitrogen solutions.

These are general product recommendations. Please consult with your NACHURS Sales Manager or agronomist for specific fertility recommendations.