

vegalab

ENVIRONMENTALLY SAFE



# NEMATODE CONTROL

NATURAL NEMATODE CONTROL  
LIQUID CONCENTRATE

<b>ACTIVE INGREDIENT:</b>		
Geraniol (CAS# 106-24-1) .....	2.5%	
<b>OTHER INGREDIENTS:*</b> .....	97.5%	
<b>TOTAL:</b> .....	100.0%	

\*Water, Lecithins (CAS# 8002-43-5), Egg Solids (CAS# 9006-50-2)

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

See Back Panel for Precautionary Statements

Manufactured by: Vegalab LLC, 1201 North Orange Street, Suite 7233, Wilmington, DE 19801-1186

**NET CONTENTS: 1 Gallon (3.79 L)**

**Stops Nematode Infestations,  
Which Enhances Plant Growth**

Nematode Control is a nematicidal agent from natural plant extracts that is used to control many species of plant parasitic nematodes, including those from the genus Meloidogyne. They exist in soil in areas with warm weather or short winters. About 2,000 plants are susceptible to infection by root-knot nematode, which causes approximately 5% of the total global crop loss. Nematode Control controls nematodes by reducing the size of root knots formed and destroying nematode eggs. It is applied to soil, where it targets plant parasitic nematodes, thereby promoting crop and root growth.

Nematode Control uses naturally extracted geraniol to destroy plant parasitic nematodes, while greatly reducing any environmental impacts. Geraniol is highly recommended for the biological control of plant parasitic nematodes.



vegalab



# NEMATODE CONTROL

NATURAL NEMATODE CONTROL

## FIRST AID

**IF SWALLOWED** • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person. **IF INHALED** • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferable mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice. **IF IN EYES** • Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice. **IF ON SKIN OR CLOTHING** • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment.

**HOT LINE NUMBER:** Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For emergency information call the National Pesticides Information Center at 1-800-858-7378, 6:30 AM to 4:30 PM Pacific time (PT), seven days a week. During other times, call the poison control center 1-800-222-1222.

- Effective against burrowing nematodes, lesion nematodes, pin nematodes, root knot nematodes, and spiral nematodes.
- Tested on cucumbers, lettuce, melons, strawberries, tomatoes (cherry), and watermelons.
- Enhances plant disease resistance, is broadly recommended, and used in various fields, such as forestry, horticulture, lawn care, and agriculture for use in all crops.

## DIRECTIONS FOR USE

Mix thoroughly and continue agitation during application. Nematode Control is only to be applied to the soil and not onto plants or leaves. If above-ground harvestable food commodities are present, spray must be directed to soil around and adjacent to plants. Nematode Control contains eggs for which the direct aerial and ground application to above-ground harvestable food commodities is prohibited. Do not apply prior to rain as this may reduce product effectiveness.

Purpose	Mix Rates	Soil Application Rates
For prevention of damage dilute 1:1,000	0.13 fl. oz. (3.8 ml)/ 1 gal. 12.8 fl. oz. (379 ml)/ 100 gal.	1.61 fl. oz. (47.8 ml)/ 1,000 ft <sup>2</sup> 67.6 fl. oz. (2.08 L)/ Acre 1.32 gal. (5 L)/ Ha
If nematode infection is already high dilute 1:500	0.26 fl. oz. (7.6 ml)/ 1 gal. 25.6 fl. oz. (758 ml)/ 100 gal.	3.23 fl. oz. (95.6 ml)/ 1,000 ft <sup>2</sup> 1.10 gal. (4.16 L)/ Acre 2.64 gal. (10 L)/ Ha

## STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

**Pesticide Storage:** Store in a cool, dry area under lock and key. Post as a pesticide storage area. Always store pesticides in the original container. Store away from food, pet food, feed, seed, fertilizers, and veterinary supplies. Place liquid formulations on lower shelves and dry formulations above.

**Pesticide Disposal:** To avoid wastes, use all material in this container by application according to label directions. If wastes cannot be avoided, offer remaining product to waste disposal facility or pesticide disposal program (often such programs are run by state or local governments or by industry).

**Container Handling:** Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill or by incineration. Do not burn unless allowed by state and local ordinances.

## ENVIRONMENTAL HAZARDS

For terrestrial uses: Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

**This product has not been registered by the United States Environmental Protection Agency. Vegalab LLC represents that this product qualifies for exemption from registration under FIFRA 25(b).**

ESL25b05 REV121815