

GROW MORE®

NUTRA-ZORB SUSPENSION CALCIUM

Neutral pH for Foliar or Soil Application

GUARANTEED ANALYSIS

Calcium (Ca)20.0%
Derived from Calcium Carbonate

CONDITION OF SALE

The seller hereby represents and warrants that the fertilizers shall be manufactured and shipped to purchaser in a form suitable for application to plants, trees and other agricultural crops. Seller further warrants that the fertilizer product shall meet the guaranteed minimum analysis as set forth on the product label. However, this warranty does not extend to the use of the product contrary to label directions and the buyer assumes the risk of any such use.

CAUTION KEEP OUT OF REACH OF CHILDREN

WARNING: This product contains chemicals known to the State of California to cause cancer and birth defects or other reproductive harm.

DENSITY: 12.5 Lbs/Gallon (1.08 Kg/Liter)

Contents sold by weight not volume.

Manufactured By
GROW MORE®
15600 New Century Drive
Gardena, CA 90248
www.growmore.com



DIRECTIONS

SOIL: APPLY 5 TO 10 GALLONS INCORPORATED INTO THE SOIL OR THROUGH IRRIGATION SYSTEM.

FOLIAR: SUGGESTED USES AS FOLLOWS

APPLES & PEARS:

Western States: Apply 2 to 8 Qts. per acre per application in pre- and/or post-bloom sprays. Repeat as needed. Do not apply during bloom period, under poor drying conditions or within 24 hours of predicted rainfall. Do not apply with other materials that may cause russetting.

APRICOTS, PRUNES, PLUMS, NECTARINES, PEACHES, CHERRIES: Apply 3 to 8 Qts. per acre per application in pre- and/or post-bloom sprays. For developing early leaf surface, use 3 to 8 Qts. per acre in pink bud stage. More than 1 application is advisable. **CAUTION:** Do not apply to fruit within 3 weeks of harvest because of visible residue.

ARTICHOKES: Apply 2 to 5 Qts. per acre per application.

ASPARAGUS: Apply 2 to 3 Qts. per acre after start of new fern growth. Repeat at intervals of 2 to 3 weeks as needed.

AVOCADOS: Apply 2 to 3 Qts. per 100 gallons and flush wet the foliage. By air or concentrate spray: use 10 to 30 Lbs. per acre per application.

BEANS, SOYBEANS, PEAS, LENTIL: Use 0.5 to 3 Qts. per acre per application. First application at bud to early bloom stage. Repeat in 2 to 3 weeks.

BLUEBERRIES, CRANBERRIES: Apply 1 to 6 Qts. per acre over new growth in Spring. Applications should not be made

after fruit starts to color.

BEETS, CARROTS, SWEET POTATOES & OTHER VEGETABLE ROOT CROPS: Apply 1 to 3 Qts. per acre per application.

CITRUS (ALL VARIETIES): By air or concentrate sprayer, apply 3 to 16 Qts. per acre per application in 15 to 100 gallons of water. By conventional ground rig: apply 1 to 3 Qts. per 100 gallons and flush wet the foliage (up to 16 Qts. per acre per application). In oil sprays, 0.5 to 1 Qts. per 100 gallons of spray mix.

GRAPES: Apply 1.5 to 5 Qts. per acre per application. First application when new shoots are in the 4 to 6-leaf stage, repeat as needed.

KIWIS: Apply 1.5 to 5 Qts. per acre per application.

LETTUCE, CELERY, COLE CROPS & LEAFY VEGETABLES: Apply 0.5 to 3 Qts. per acre per application. Make several applications at intervals of 2 to 3 weeks starting at thinning period.

MELONS, CUCUMBERS & OTHER CUCURBITAE: Apply 0.5 to 3 Qts. per acre per application. Make 2 or more applications at intervals of 2 to 3 weeks starting just prior to bloom.

MINT: Apply 1.5 to 5 Qts. per acre per application. First application at 6 inches of new growth, repeat 2 to 3 weeks later or as needed.

NUT CROPS (ALMONDS, WALNUTS, PECANS, PISTACHIOS, FILBERTS): Apply 3 to 10 Qts. per acre per application, pre-bloom and in cover sprays.

OLIVES: Apply 5 to 10 Qts. per acre per application. Apply 15 to 20 days after thinning, repeat 3 to 4 weeks later. If crop is not thinned, apply 1 to 2 weeks after full bloom, repeat 3 to 4

weeks later.

PEANUTS: Apply 1.5 to 10 Qts. per acre per application. First application at early-bloom stage, repeat at intervals of 2 to 3 weeks as needed.

POTATOES: Apply 1.5 to 5 Qts. per acre per application. First application when plants are 4 to 12 inches high. Second application 3 weeks later, repeat as needed.

STRAWBERRIES: Apply 1.5 to 3 Qts. per acre per application. Do not apply after fruit starts to color as a visible residue may remain.

SUGAR BEETS: Apply 1.5 to 5 Qts. per acre per application. First application when plants are 6 to 8 inches across. Second application 3 to 4 weeks later, repeat as needed.

TOMATOES, PEPPERS: Apply 1 to 3 Qts. per acre per application in enough water to give even distribution of materials. One to three applications are advised at 3 to 4-week intervals, starting in period prior to bloom.

DILUTION FOR AIR OR CONCENTRATE SPRAYS: When total gallonage applied per acre is 25 gallons or less, NUTRA-ZORB may be used at the rate of 1/2 Pint (8 Fl. oz.) per gallon of water, but DO NOT EXCEED THE MAXIMUM RECOMMENDED RATE PER ACRE.

COMPATIBILITY: NUTRA-ZORB is compatible with most commonly used wettable powder fungicides, insecticides and miticides.

SPREADER: The addition of 1/2 pint to 1 quart per 100 gallons (1/2 to 2 pints per acre by air or concentrate rig) as a spreader and acidifier will improve results on all crops.

2.5 GALLON (9.46 LITERS)