

NUTRAMIX™

0-28*-25 FOLIAR NUTRIENT | SOIL NUTRIENT

12.3 LBS. PER GAL @ 68 ° F
1376 GRAMS PER LITER @ 20 ° C

GUARANTEED ANALYSIS

TOTAL AVAILABLE PHOSPHORIC ACID (P₂O₅) AOAC METHOD 960.02). 28.0%
SOLUBLE POTASH (K₂O)..... 25.0%

DERIVED FROM POTASSIUM HYDROXIDE AND PHOSPHOROUS ACID

DO NOT APPLY WITH COPPER

KEEP OUT OF REACH OF CHILDREN

WARRANTY: WTB TECHNOLOGY makes no warranty, express or implied, including the warranties of merchantability and/or fitness for any particular purpose, concerning this material, except those which are contained on the WTB TECHNOLOGY label attached to the product container.

Information regarding the contents and levels of metals in this product is available by calling 1-800-542-6664

PRODUCT INFORMATION

NUTRAMIX PK is a highly concentrated nutritional solution. It is designed to meet the need for a soluble liquid minor element spray containing certain nutrients proven to be of value when applied as a properly timed foliage spray. NUTRAMIX PK becomes generally more available for plant food and non-phytotoxic when used as directed. It is absorbed through the leafy tissue of the plant and can be translocated within the plant. It is Phosphorous acid products that are intended to be used as a supplemental fertilizer treatment.

NUTRAMIX PK nutrients can be applied by conventional ground rig, concentrate sprayer, and by air. For improved, leaf absorption add 1-4 qts. (.95-3.80 liters) per acre. NUTRAMIX PK also has a buffering effect on hard water.

USERS CAUTION: Harmful if swallowed. Avoid contact with skin and eyes. In case of contact, flush with water.

APPLICATION CAUTION: A compatibility test is recommended if mixing NUTRAMIX PK in a tank mix. User assumes full responsibility for product, equipment and plant compatibility. Upon foliar application, the phosphite ions are taken up directly by the plant foliage and may undergo a degree of conversion to phosphate ions, or be utilized directly by the plants, as phosphite ions. As a soil application to annual crops, a lesser response from the initial crop, with a corresponding superior response from succeeding crops, may be observed. In addition, placement close to seeds or root zones may be injurious to crops. The effect may be aggravated by a soil pH below 6.5.

SOIL APPLICATION: NUTRAMIX PK can be applied to soil through various types of irrigation and application systems. The application rate is 32 to 64 ounces per acre. Do not exceed 2 gallons per 100 gallons of water or 2% volume per application.

* As a soil application to annual crops, a lesser response from the initial crop, with a corresponding superior response from succeeding crops, may be observed. In addition, placement close to seeds or root zones may be injurious to crops. The effect may be aggravated by a soil pH below 6.5

FOLIAR APPLICATION: NUTRAMIX PK may be applied by air or ground in concentrate or dilute sprays. For aerial application, apply 1 to 4 pints per acre in a minimum of 20 gallons of water per acre. For ground application, apply 1 to 4 pints per acre, and do not exceed a product concentration of 2% volume (for each pint of 0-28*-25, add 6 1/4 gallons of water). * Upon foliar application, the phosphite ions are taken up directly by the plant foliage and may undergo a degree of conversion to phosphate ions, or be utilized directly by the plants, as phosphite ions.

APPLICATION RATES

AVOCADOS, WALNUTS, ALMONDS, PISTACHIOS, FILBERTS, AND PECANS - 1-4 qts. (0.95-3.80 liters) per acre three times each season starting at first full leaf, early nut deve lopment, and three to four weeks prior to hull split.

BLACKBERRIES, BOYSENBERRIES, RASPBERRIES, STRAWBERRIES and other small fruits - 1-4 qts. (0.95-3.80 liters) per acre after transplant or thinning. Repeat every 10-15 days.

BROCCOLI, BRUSSEL SPROUTS, CAULIFLOWER, CABBAGE - 1-4 qts. (0.95-3.80 liters) per acre after transplant or thinning. Repeat every 10-15 days.

CARROTS, CUCUMBERS & SQUASH - 1-4 qts. (0.95-3.80 liters) per acre after transplant or thinning. Repeat every 10-15 days.

LETTUCE, CELERY AND OTHER LEAFY CROPS - 1-4 qts. (0.95-3.80 liters) per acre after transplant or thinning. Repeat every 10-15 days.

MELONS -1-4 qts. (0.95-3.80 liters) per acre after transplant or thinning. Repeat every 10-15 days.

ONIONS, LEEKS, PEAS, BEANS, PEPPERS AND GARLIC - 1-4 qts. (0.95-3.80 liters) per acre after transplant or thinning. Repeat every 10-15 days.

SPINACH, SUGAR BEETS, TOMATOES, EGGPLANT - 1-4 qts. (.95-3.80 liters) per acre after transplant or thinning. Repeat every 10-15 days.

POTATOES - 1-4 qts. (.95-3.80 liters) per acre three to four times during season. The first at early emergence, three to four weeks later and pre-bloom.

VINE CROPS -

GRAPES - 1-4 qts. (.95 - 3.80 liters) per acre three to four times each season 1-4 qts. (.95 - 3.80 liters) per acre at first full leaf, then 1-4 qts. .95 - 3.80 liters) per acre at post bloom berry size and three to four weeks prior to harvest.

TREE CROPS

APPLES, NECTARINES, PEACHES, PEARS, PLUMS & CITRUS - 1-4 qts. (.95 - 3.80 liters) per acre three to four times each season starting at three quarter leaf then at early fruit size, fruit midsize and three to four weeks prior to harvest.

NUTRAMIX IS A REGISTERED TRADEMARK

Production #: 707 Date: 07/10/2020



MANUFACTURED BY
WTB TECHNOLOGY CORPORATION
245 Industrial Street • Bakersfield • California 93307
(661) 327-9649 | Fax (661) 327-1740 | E-mail: info@westernnutrientscorp.com