

**FIFRA SECTION 24(C) SPECIAL LOCAL NEED LABEL
FOR DISTRIBUTION AND USE ONLY IN THE STATE OF OREGON**

PYRIDABEN	GROUP	21A	ACARICIDE
-----------	-------	------------	-----------



EPA Reg. No 81880-28-10163 / EPA SLN NO. OR-170008

For Control of Pear Psylla Nymphs on Pears

This label is valid until December 31, 2027, or until otherwise amended, withdrawn, or canceled or suspended.

ACTIVE INGREDIENT: PYRIDABEN

% By Wt.

[2-tert-butyl-5-(4-tert-butylbenzylthio)-4-chloropyridazin-3(2H)-one] 42.47%

OTHER INGREDIENTS: 57.53%

Total 100.00%

Contains 3.75 pounds active ingredient per gallon

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high-water mark. Keep out of lakes, ponds, or streams. Do not contaminate water by cleaning of equipment or disposal of equipment washwaters. Do not apply when weather conditions favor drift from target area. Drift or runoff from treated areas may be hazardous to fish in adjacent sites. This product is highly toxic to bees. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively foraging the treatment area.

POLLINATOR PROTECTION: Nexter SC Miticide/Insecticide may only be applied between late evening and early morning.

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This FIFRA Section 24(c) Special Local Need (SLN) label and the main EPA-registered label for Nexter SC must be in the possession of the user at the time of pesticide application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on this SLN label and the main EPA-registered label.
- REI: Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours. Exception: the REI is 2 days for treated pears when hand thinning.

CROP	RATE	PEST	COMMENTS
Pears (including oriental)	17 – 25 fl. oz. / acre (0.5-0.73 lb ai/acre)	Pear Psylla	Apply Nexter SC using conventional dilute or concentrate ground sprayers calibrated to deliver enough water to ensure thorough coverage of foliage. Spray volume will vary with the size and number of trees per acre and foliage density. Nexter SC must be applied to each row for maximum coverage. Pear psylla numbers can increase rapidly; begin application before damaging numbers are reached. Applications may be made early from cluster bud through petal fall to control eggs and early pear psylla instars. Nexter SC is also effective when applied after petal fall as psylla populations begin to build.
RESTRICTIONS: <ul style="list-style-type: none">• Do not apply within 28 days of harvest (PHI).• Do not make more than 1 application per year.• Do not apply using aerial application.• For use under this SLN label do not apply this product through any type of irrigation system.• Applicators applying this product by airblast application at the maximum labeled rate must wear chemical-resistant headgear, unless applying in an enclosed cab. This requirement is in addition to the Personal Protective Equipment (PPE) requirements on the main EPA-registered label.• Drift: Do not apply within 110 feet upwind of surface water. Follow all applicable directions, restrictions and precautions related to spray drift management on the main EPA- registered label.			

Nexter® is a registered trademark of Nissan Chemical Corporation

24(c) Registrant: Canyon Group, LLC
c/o Gowan Company
P.O. Box 5569
Yuma, AZ 85366-5569

Distributor: Gowan Company, LLC
P.O. Box 5569
Yuma, AZ 85366-5569