



Bayer CropScience LP
 800 N.Lindbergh Blvd.
 St. Louis, MO 63167
 1-866-99BAYER (1-866-992-2937)

Nortron® SC Herbicide

EPA Reg. No. 264-613

For selective control of weeds in garden beets.

EPA SLN No. NY-120014

FIFRA Section 24(c) Special Local Need (SLN)

**FOR DISTRIBUTION AND USE WITHIN THE STATE OF NEW YORK
 EXCEPT NASSAU AND SUFFOLK COUNTIES**

THIS LABEL IS VALID UNTIL **DECEMBER 31, 2023** UNLESS OTHERWISE AMENDED, WITHDRAWN, CANCELED, OR SUSPENDED.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its label. This label must be in the possession of the user at the time of application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered Section 3 label.

Failure to follow the Directions for Use and Precautions on this label may result in poor product performance, crop injury, or illegal residues.

Apply the rate of Nortron® SC Herbicide recommended below using ground equipment in 10 to 40 gallons of water before weeds emerge as a preemergent, preplant incorporated or post-emergence treatment (see Chart 1). Nortron® SC has limited activity on emerged weeds. (See complete Nortron® SC Herbicide label for recommendations on soil preparation, spray equipment, incorporation equipment, and cultural practices. User must comply with rotational and plantback restrictions specified in the General Use Precautions).

CHART 1. Dosage Chart for of Nortron® SC Herbicide alone.

BEETS, TABLE (GARDEN)

DOSAGE FOR BROADCAST APPLICATIONS

Table Beet Stage	NORTRON® SC HERBICIDE FLUID OUNCES/ACRE BROADCAST
Preemergence	60
Postemergence	
2-Leaf	5.25
4-Leaf	5.25
6-Leaf to 8-Leaf	10.5

Do not exceed 96 fluid ounces (0.75 gallon) of product in a single growing season (3 lbs ai/A per season). Do not make more than 2 applications of Nortron® SC Herbicide per acre per season. See *Use Restrictions* on the EPA Registered label for additional information on proper use.

PRECAUTIONS: Higher rates of Nortron® SC applied after planting may result in crop injury. Crop injury may result under unfavorable environmental conditions if the crop is under stress from one or more of the following conditions:

- Rapid climatic changes from cool, overcast days, to hot (80°F or over), bright days. When the air temperature is, or is likely to be, above 80°F on the day of spraying, application should be made in the evening when the temperature is lower.
- Frost within seven days following treatment
- Drought conditions

USE RESTRICTIONS

- Nortron® SC Herbicide or tank mixes must be used for registered purposes and at specified rates only. (DO NOT OVERTREAT.)
- Do not graze or feed any crop treated with Nortron SC Herbicide within 60 days of the application.
- Do not apply this product through any type of irrigation system.
- Do not allow spray mixture to stand in tank overnight. Flush and drain spray equipment after each day's use.
- Do not use Nortron® SC Herbicide on muck or peat soils.
- Store unused spray mixture in tightly-sealed containers and protect from frost.
- This label must be in the possession of the user at the time of pesticide application.
- Distribution, use and sale of Nortron® SC Herbicide are prohibited in Nassau & Suffolk counties.

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)

As with any plant-protection product, always read and follow label directions.