



Bayer CropScience LP
 P.O. Box 12014
 2 T.W. Alexander Drive
 Research Triangle Park, North Carolina 27709
 1-866-99BAYER (1-866-992-2937)

Nortron® SC Herbicide

For Control of Ladysthumb, Pale Smartweed, Wild Buckwheat, and Prostrate Knotweed in Table Beets and Swiss Chard Grown for Seed.

EPA Reg. No. 264-613

SLN. No. WA-010002

FIFRA 24(c) Special Local Need

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF WASHINGTON

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. This label must be in the possession of the user at the time of pesticide application. All applicable directions, restrictions and precautions on the EPA registered label are to be followed. This label must be in the position of the user at the time of application.

This label for Nortron SC Herbicide expires and must not be distributed or used in accordance with this SLN registration after December 31, 2019.

DIRECTIONS FOR USE

Failure to follow the Directions for Use and Precautions on this label may result in poor product performance, crop injury, or illegal residues.

Apply the rate of Nortron® SC Herbicide recommended below using ground equipment in 10 to 40 gallons of water before weeds emerge as a preemergent, preplant incorporated or post-emergence treatment (see Chart 1). Nortron SC has limited activity on emerged weeds. (See complete Nortron® SC Herbicide label for recommendations on soil preparation, spray equipment, incorporation equipment, and cultural practices. User must comply with rotational and plantback restrictions specified in the General Use Precautions.)

CHART 1. Dosage Chart for Nortron® SC Herbicide

		Pints/Acre/Application
Direct Seeded	Pre-plant Incorporation	3.0 pints
	Preemergence	4.0 pints
Stecklings (over-wintered table beet roots) and seedlings (Swiss chard and table beets)	Post-emergence	Apply up to 1-3 pints of Nortron® SC. Use the lower rate range on light soils and higher rate range on heavier soils. Apply up to 2 pints when the crop is in the 4 to 5 leaf stage of growth and up to 3 pints when the crop is in the 6 to 8 leaf stage of growth.
Stecklings (over-wintered table beet roots and seedlings (Swiss chard and table beets)	Pre-plant incorporated in heavy soils.*	6.0 pints

***Note: The 6 pint rate is for pre-plant incorporated only.**

RESTRICTIONS:

- Allow a minimum of 28 days (4 weeks) between applications of Nortron® SC Herbicide.
- Do not make more than 2 applications Nortron® SC herbicide per acre per year.
- Do not apply Nortron® SC after the crop begins to form buds.
- For use under this SLN label, do not apply this product through any type of irrigation system.
- All table beet and Swiss chard seed screenings shall be disposed of in such a way that they cannot be distributed or used for food or feed. The seed conditioner shall keep records of screening disposal for three years from the date of disposal and shall furnish the records to the WSDA forthwith upon request. Disposal records shall consist of documentation from a controlled dumpsite, incinerator, or other equivalent disposal site and shall show the lot numbers, amount of material disposed of, it's grower(s), and the date of disposal. No portion of the table beet and Swiss chard seed plant, including but not limited to green chop, hay, pellets, meal, whole seed, cracked seed, roots, bulbs, leaves and seed screenings may be used or distributed for food or feed purposes. Table beet and Swiss chard seed shall bear a tag or container label, which forbids use of the seed for human consumption or animal feed. Table beet and Swiss chard seed may not be distributed for human consumption or animal feed.
- Producers of table beet or Swiss chard seed who use this product are required to inform, in writing, conditioners receiving seed produced on fields treated with this product. A copy of this labeling is required to be provided to the conditioners by the producer. At the processing plant, processed seed must be labeled "Not for human or animal consumption". The processor must dispose of all seed screenings in such a way that they cannot be distributed or used for food or feed.

WSDA Container Disposal Guidance: Pesticide containers must be properly cleaned prior to disposal. The best time to clean empty pesticide containers is during mixing and loading, because residue can be difficult to remove after it dries. Triple rinse (or pressure rinse) the pesticide container, empty all pesticide rinse water into the spray tank, and apply to a labeled crop or site. Recycling cleaned containers is the best method of container disposal. Information regarding the recycling of empty and cleaned plastic pesticide containers in Washington is available on the WSDA Waste Pesticide Program web site at <http://agr.wa.gov/PestFert/Pesticides/WastePesticide.aspx>. Cleaned containers may also be disposed of in a sanitary landfill, if permitted by the county. Burning is not a legal method of container disposal in Washington.

PRECAUTIONS: Higher rates of Nortron® SC applied after planting may result in crop injury. Crop injury may result under unfavorable environmental conditions if the crop is under stress from one or more of the following conditions:

- Rapid climatic changes from cool, overcast days, to hot (80°F or over), bright days. When the air temperature is, or is likely to be, above 80°F on the day of spraying, application should be made in the evening when the temperature is lower.
- Frost within seven days following treatment.
- Drought conditions.

**Bayer CropScience Special Conditions and Disclaimer for Use of Nortron SC Herbicide
on Table Beets or Swiss Chard grown for seed.**

Bayer CropScience intends that this Section 24(c) label be distributed only by the Puget Sound Seed Growers Association only to end users and/or growers who agree in writing to the terms and conditions required by the Puget Sound Seed Growers Association including a waiver and release from all liability and indemnification by the user and/or grower of Bayer CropScience, Puget Sound Seed Growers Association, and others for failure to perform and crop damage from the use of Nortron® SC Herbicide on table beet and Swiss chard grown for seed. If such terms and conditions are unacceptable, return Nortron® SC Herbicide at once unopened.

This product when used on table beet and Swiss chard grown for seed may lead to crop injury, loss or damage. Bayer CropScience recommends that the user and/or grower test this product in order to determine its suitability for such intended use. The Puget Sound Seed Growers Association and Bayer CropScience make this product available to the user and/or grower solely to the extent the benefit and utility, in the sole opinion of the user and/or grower, outweigh the extent of potential injury associated with the use of this product. The decision to use or not to use Nortron® SC Herbicide must be made by each individual Nortron® SC Herbicide user and/or grower on the basis of possible crop injury from Nortron® SC Herbicide, the severity of weed infestation, the cost of alternative (pest) controls, and other factors. Bayer CropScience intends that because of the risk of failure to perform or crop damage that all such use is at the user's and/or grower's risk, to the extent consistent with applicable law.

This Special Conditions and Disclaimer statement is required by Bayer CropScience and is not required or enforced by the USEPA or the WSDA.

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577
For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)

24(c) Registrant:

Bayer CropScience LP

P.O. Box 12014, 2 T.W. Alexander Drive

Research Triangle Park, North Carolina 27709

Nortron is a registered trademark of Bayer.