

FIFRA Section 24(c) Special Local Need (SLN) Label

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF OREGON

Oberon[®] 4 SC Insecticide/Miticide

EPA Reg. No. 264-850

EPA SLN. No. OR-200013

For Control of Twospotted Spider Mite in Mint

This label is valid until December 31, 2025 or until otherwise amended, withdrawn, canceled, or suspended.

ACTIVE INGREDIENT: Spiromesifen:

2-oxo-3-(2,4,6-trimethylphenyl)-1-oxaspiro[4.4]non-3-en-4-yl 3,3- dimethylbutanoate..... 45.20%

OTHER INGREDIENTS: 54.80%

TOTAL: 100.00%

Oberon contains 4 pounds of spiromesifen per US gallon (480 grams per liter).

KEEP OUT OF REACH OF CHILDREN

CAUTION

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates. Do not contaminate surface water through spray drift. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwater.

Surface Water Advisory

This product may contaminate water through runoff or drift of spray in wind. Poorly draining soils and soils with shallow water tables are more prone to produce runoff that contains this product. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from rainfall runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This Special Local Need (SLN) label and the EPA-registered Oberon 4 SC Insecticide/Miticide Section 3 product container label must be in the possession of the user at the time of pesticide application.

Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label for Oberon 4 SC Insecticide/Miticide (EPA Reg. No. 264-850) and this SLN label.

MINT

| Pests Controlled | RATE PER APPLICATION fluid ounces/acre | Use Directions |
|------------------------|---|--|
| Twospotted spider mite | 4.0 - 8.0 | Make a foliar application at the onset of infestation, prior to the establishment of a damaging population. Repeat as necessary but no closer than a 7-day interval. Application may be made by ground, by air, or by chemigation (refer to Restrictions/Precautions). Please refer to the OBERON 4 SC Section 3 label for additional information. |

RESTRICTIONS/PRECAUTIONS

Pre-harvest Interval (PHI): 7 days.

Maximum OBERON 4 SC INSECTICIDE/MITICIDE allowed per 7-day interval: 8.0 fl oz per acre (0.25 lb ai/A).

Maximum OBERON 4 SC INSECTICIDE/MITICIDE allowed per crop season: 24.0 fl oz per acre (0.75 lb ai/A).

Maximum number of applications per crop season: 3.

Minimum application volume: 10.0 GPA – ground, 5.0 GPA – aerial application.

Do not apply during mint flowering. Do not apply when bees are foraging on mint.

This product may be applied through irrigation systems. Refer to the EPA registered label for chemigation directions.

Types of Irrigation Systems: Apply OBERON 4 SC INSECTICIDE/MITICIDE only through sprinkler, including center pivot, lateral move, side roll, or overhead solid set irrigation systems. Do not apply OBERON 4 SC INSECTICIDE/MITICIDE through any other type of irrigation system.

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577.

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937).

As with any crop-protection product, always read and follow label directions.

For additional information call toll-free 1-866-99BAYER (1-866-992-2937).

10/02/2020

Section 24c SLN Registrant:

Bayer CropScience, LP
800 N. Lindbergh Blvd.
St. Louis, MO 63167