

PRODUCT INFORMATION BULLETIN

FIFRA Section 2(ee) Recommendation
Expiration Date: October 1, 2022

FOR DISTRIBUTION AND USE ONLY IN THE STATE OF TEXAS

POTASSIUM PHOSPHITE	GROUP	33	FUNGICIDE
TEBUCONAZOLE	GROUP	3	FUNGICIDE
AZOXYSTROBIN	GROUP	11	FUNGICIDE



A systemic fungicide for the control of corn and soybean diseases

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or accepted by the U.S. Environmental Protection Agency. This product bulletin should be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product inconsistent with its labeling.

<u>CROP: CORN* (Field corn)</u>		
<u>DISEASE</u>	<u>RATE</u>	<u>INSTRUCTIONS</u>
Rust (<i>Puccinia</i> spp.) ----- Northern leaf blight (<i>Exserohilum turcicum</i>) Southern leaf blight (<i>Bipolaris maydis</i>) Northern leaf Spot (<i>Bipolaris zeicola</i>) Gray leaf spot (<i>Cercospora zeae-maydis</i>) Tar Spot (<i>Phyllachora maydis</i>)	2-3 pints/acre ----- 2.25-3 pints/acre	Apply the higher specified rate (3 pts/acre) when conditions favor severe disease development. Apply preventatively when weather favors disease development. Repeat on a 7-14 day interval.
Specific Use Restrictions: 1) Do not make more than 2 applications per year. 2) Do not apply more than 0.45 pounds of ai tebuconazole-containing products/A/year. 3) Do not apply more than 2.0 pounds of ai azoxystrobin-containing products/A/year 4) Do not apply more than 6 pints per acre of product per year. 5) Do not apply within 7 days of harvest (PHI=7 days). * Not for use in California, New York, and Washington.		

<u>CROP: SOYBEAN</u>		
<u>DISEASE</u>	<u>RATE</u>	<u>INSTRUCTIONS</u>
Soybean rust (<i>Phakopsora pachyrizi</i>) ----- Downy mildew (<i>Peronospora manshurica</i>), Powdery mildew (<i>Microsphaera diffusa</i>), Frogeye leafspot (<i>Cercospora sojina</i>), Aerial blight (<i>Rhizoctonia solani</i>), Brown Leaf spot (<i>Septoria glycines</i>)	2 pints ----- 2.25 - 2.5 pints/acre	For rust, apply preventively when weather conditions favor disease development. Repeat applications on a 7-14 day interval. Apply the higher specified rate (2.5 pts/acre) when conditions favor severe disease development Make applications between R1 and R5 (flowering through visible seed in the pod in the upper 4 nodes of the plant). Make a second application 21 days later if necessary. Use spray volumes of 2 gallons or more by air, and 10 gallons or more by ground. To prevent mixing issues, add recommended surfactant. Keep under constant agitation during spraying. Use adjuvants at their lowest labeled rate.
Specific Use Restrictions: <ol style="list-style-type: none"> 1) Do not make more than 2 applications per year. 2) Do not apply more than 0.225 pounds of tebuconazole-containing products/A/year. 3) Do not apply more than 1.5 pounds of ai azoxystrobin-containing products/A/year. 4) Do not apply more than 4 pints per acre of product per year. 5) Do not apply within 21 days of harvest (PHI=21 days). <p>* Not for use in California, New York, and Washington.</p>		

EPA Reg. No. 42519-40-5905

Manufactured for
HELENA AGRI-ENTERPRISES, LLC
225 SCHILLING BOULEVARD, SUITE 300
COLLIERVILLE, TN 38017