

---

## SUPPLEMENTAL LABELING

---

**Syngenta Crop Protection, LLC**

P. O. Box 18300

Greensboro, North Carolina 27419-8300

SCP 71512A-S4 0516

**GROUP 29 FUNGICIDE****SUPPLEMENTAL DIRECTIONS FOR THE USE OF  
OMEGA® 500F  
(EPA Reg. No. 71512-1-100)****FOR CONTROL OF DISEASES ON MELONS**

This Supplemental Label Expires on May 30, 2019 and must not be used or distributed after this date.

**Directions for Use:**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DISEASES	PRODUCT RATE PER ACRE
<b>CROP: CUCURBIT VEGETABLES, MELON SUBGROUP 9A</b>	
Phytophthora blight <i>Phytophthora capsici</i> Downy mildew <i>Pseudoperonospora cubensis</i> Gummy stem blight <i>Didymella bryoniae</i> Alternaria leaf spot <i>Alternaria cucumerina</i>	0.75 to 1.5 pints
<b>APPLICATION DIRECTIONS:</b>	
<p>For Phytophthora blight control the first application may be made at 1.5 pt/A as a banded soil drench at transplant or when the plants have the first true leaves. Subsequent foliar applications for Phytophthora blight and downy mildew should be made at 0.75 to 1 pt/A on a 7 – 10-day interval beginning when disease first appears or when conditions are favorable for disease development. Use the low rate when conditions are favorable for disease development or when disease pressure is low to moderate. Use sufficient water to provide coverage of the foliage. For Phytophthora blight and gummy stem blight, applications should be directed to provide coverage of the lower stem area.</p> <p>Use the low rate and longest interval for preventative applications and when disease pressure is low. Increase the rate and decrease the interval as disease pressure increases. For high disease pressure use the 1.5 pint rate on a weekly interval.</p> <p>DO NOT apply more than 9 pints of Omega 500F (4.69 lbs ai) per acre per year. DO NOT apply within 30 days of harvest. Restricted Entry Interval (REI) = 12 hours DO NOT apply this product with mechanically pressurized handgun equipment.</p> <p>Omega 500F may be applied through sprinkler system irrigation equipment on cucurbits. See irrigation use directions elsewhere on the Omega 500F label. Includes all members of the Cucurbit Vegetables, Melon Crop Subgroup 9A, such as: Citron melon;</p>	

Muskmelon, including hybrids and/or varieties of *Cucumis melo* (including true cantaloupe, cantaloupe, casaba, Santa Claus melon, Crenshaw melon, honeydew melon, honey balls, Persian melon, golden pershaw melon, mango melon, pineapple melon, snake melon); and watermelon, *Citrullus spp.*; including hybrids and/or varieties of these.

**NOTE:** Follow all applicable directions, restrictions, and precautions on the **Omega 500F** label.

**NOTE:** This labeling must be in the possession of the user at the time of pesticide application.

©2017 Syngenta

**OMEGA 500F - SCP 71512A-S4 0516 – dk – 1/23/17**