
SUPPLEMENTAL LABELING

Syngenta Crop Protection, LLC

P. O. Box 18300

Greensboro, North Carolina 27419-8300

SCP 71512A-S5 0516

GROUP	29	FUNGICIDE
--------------	-----------	------------------

SUPPLEMENTAL DIRECTIONS FOR THE USE OF
OMEGA® 500F
(EPA Reg. No. 71512-1-100)

FOR CONTROL OF DISEASES ON TUBEROUS AND CORM VEGETABLES (SUBGROUP 1C)This Supplemental Label Expires on May 30, 2019 and must not be used or distributed after this date.**Directions for Use:**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DISEASES	PRODUCT RATE PER ACRE
CROP ROOT AND TUBER VEGETABLES Tuberous and Corm Subgroup 1C	
Late blight (<i>Phytophthora infestans</i>)	5.5 to 8 fl oz
White mold (<i>Sclerotinia sclerotiorum</i>)	
APPLICATION DIRECTIONS: For late blight and white mold control, begin applications when the plants are 6 to 8 inches tall or when conditions favor disease development. Repeat applications at intervals of 7 to 10 days. When white mold pressure is low to moderate, use 5 ½ fluid ounces. When conditions favor moderate to high white mold pressure increase the rate to 8 fluid ounces.	
Restrictions DO NOT apply more than 3.5 pints of Omega 500F (1.82 lbs ai) per acre per year. DO NOT apply within 14 days of harvest DO NOT apply this product with mechanically pressurized handgun equipment. Restricted Entry Interval, REI = 12 hours Omega 500F may be applied by aerial application (except in the State of New York) or through sprinkler system irrigation equipment on potatoes. See irrigation use directions on full label. Includes all members of Root and Tuber Vegetables, Tuberous and Corm Crop Subgroup 1C: Arracacha; arrowroot; artichoke, Chinese; artichoke, Jerusalem; canna, edible; cassava, bitter and sweet; chayote (root); chufa; dasheen (taro); ginger; leren; potato; taniel; turmeric; yam bean; yam, true; cultivars, varieties and/or hybrids of these.	

NOTE: Follow all applicable directions, restrictions, and precautions on the **Omega 500F** label.
NOTE: This labeling must be in the possession of the user at the time of pesticide application.

©2017 Syngenta

OMEGA 500F - SCP 71512A-S5 0516 – dk – 1/23/17