
SUPPLEMENTAL LABELING

Syngenta Crop Protection, LLC

P. O. Box 18300

Greensboro, North Carolina 27419-8300

SCP 71512A-S12 1021

GROUP	29	FUNGICIDE
-------	-----------	-----------

SUPPLEMENTAL DIRECTIONS FOR THE USE OF

OMEGA 500F
(EPA Reg. No. 71512-1-100)

FOR CONTROL OF DISEASES ON TOMATO FRUITING VEGETABLES

This Supplemental Label expires on October 31, 2024 and must not be used or distributed after this date.

Directions for Use:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

CROP: TOMATO, FRUITING VEGETABLE, SUBGROUP 8-10A; INCLUDES ALL MEMBERS OF FRUITING VEGETABLE, TOMATO SUBGROUP 8-10A, INCLUDING: BUSH TOMATO, COCONA; CURRANT TOMATO; GARDEN HUCKLEBERRY; GOJI BERRY; GROUNDCHERRY; NARANJILLA; SUNBERRY; TOMATILLO; TOMATO; TREE TOMATO; CULTIVARS, VARIETIES, AND/OR HYBRIDS OF THESE.

DISEASES	RATE PER ACRE
Southern blight (<i>Athelia rolfsii</i>)	16 to 24 fl. oz. (0.521 to 0.782 lbs. a.i.)

APPLICATION DIRECTIONS

To control Southern blight on tomatoes, apply the initial application as a soil directed spray at or within the day of transplanting at 24 fl. oz. (0.782 lbs. a.i.)/A. Direct the spray towards the soil and base of the plant along each side of the row. Irrigate within one hour of the application with approximately ½ to 1 inch of water to properly move the product into the root zone. Follow the soil directed spray at transplant with up to five foliar applications at 7- to 14-day retreatment intervals.

For foliar applications use the low rate and longest interval for preventative applications and when disease pressure is low. For moderate disease pressure use the 16 fl. oz. (0.521 lbs. a.i.) rate on a weekly interval. For high disease pressure use the 24 fl. oz. (0.782 lbs. a.i.) rate on a weekly interval. Use adequate water to provide coverage of foliage.

Restrictions

DO NOT apply more than 144 fl. oz. of Omega 500F (4.69 lbs. a.i.) per acre per year.

DO NOT make more than 6 applications to tomatoes at the 24 fl. oz./A rate/year.

DO NOT apply to tomatoes within 7 days of harvest (PHI = 7 days).

Restricted Entry Interval (REI) = 12 hours.

The maximum single use rate is 24 fl. oz. (0.782 lbs. a.i.)/A with the shortest RTI of 7 days.

OMEGA 500F may be applied through sprinkler system irrigation equipment on fruiting vegetables. See calibration directions elsewhere on the Omega 500F label.

NOTE: Follow all applicable directions, restrictions, and precautions on the **Omega 500F** label.

NOTE: This labeling must be in the possession of the user at the time of pesticide application.

©2021 Syngenta

OMEGA 500F - SCP 71512A-S12 1021 – dk – 3/8/22