

OrganOmex™



Achieve 3-1-3



Guaranteed Analysis

Total Nitrogen (N).....	3.00%
3% Water Soluble Nitrogen	
Available Phosphate (P ₂ O ₅)	1.00%
Soluble Potash (K ₂ O)	3.00%

Derived from: Fermented Guayule Extract, Soy Protein Hydrolysate, Sulfate of Potash, Hydrolyzed Fish and Phosphoric Acid (as stabilizer)
F2737

KEEP OUT OF REACH OF CHILDREN

Hazard Statements: This product is not classified as a hazardous chemical under OSHA Hazard Communication Standard (29 CFR 1910.1200 (HCS))

Precautionary Statements: Wash face, hands and any exposed skin thoroughly after handling. Do not breathe spray. Wear protective gloves, clothing and eye protection. If swallowed, rinse mouth. Do not induce vomiting. Dispose of contents/container according to regulations.

FOR AGRICULTURAL USE ONLY

GUARANTEED BY:

OMEX® AGRIFLUIDS, INC. • PO BOX 815 • SELMA, CA 93662



DENSITY: 11.09 LBS/GALLON @ 68° F

LIQUID CONTENT 1 US GALLONS (3.78 L)

LIQUID CONTENT 2.5 US GALLONS (9.5 L)

LIQUID CONTENT 275 US GALLONS (1,040 L)



NET WEIGHT: 11.09 LBS (5.03 kg)

NET WEIGHT: 27.73 LBS (12.57 kg)

NET WEIGHT: 3,060 LBS (1,388 kg)

GENERAL INFORMATION

OrganOmex™ Achieve 3-1-3 is designed to deliver nitrogen, phosphorus and potassium efficiently to optimize nutrient levels to enhance rooting, blossoms, fruit yields and quality. Use OrganOmex™ Achieve 3-1-3 in addition to a primary fertility program.

DIRECTIONS FOR USE For Commercial Application Only.

ALFALFA: Summer Cuttings: Apply 1/2-pint per acre (8oz) in spring and 1/2 pint (8oz) every other cutting to a maximum of 2 pints for crop year. Apply as a fertigation or foliar application after cutting and removal of cut hay.

CORN and Sorghum: Apply 2 oz per acre In-Furrow followed by . Apply 6 oz per acre 2nd – 3rd Node. For single application, apply 8 oz at 2nd - 3rd Node. Late season, apply ½ pint per acre (8 oz) at Flag/leaf or pre-tassel thru fertigation or foliar. Total application should not exceed 1-pint per acre per crop cycle.

CEREAL GRAINS, such as Barley, Buckwheat, Canola, Hops, Millet, Oats, Popping Rice, Rye Sesame, Wheat, and Wild Rice: Apply ½-pint per acre (8 oz) as a fertigation or foliar application at tillering (3-6 leaf stage). Total applications should not exceed 1 pint per acre per crop cycle.

CITRUS, such as Calmondin, Citrus Hybrids, Grapefruit, Kumquat, Lemon, Lime, Mandarin, Orange (sour and sweet), Pummelo, and Tangerine: Apply 1/2 pint per acre (8oz) as a fertigation or foliar application at pre-bloom or one time between full bloom and petal fall. Repeat application during the early sizing period 2 weeks after petal fall. **FOR YOUNG TREES:** Apply ½ pint per (8 oz) acre in spring and ½ pint per acre in fall for growth. **COTTON:**

FULL SEASON, IRRIGATED COTTON. Apply 1-pint per acre (16 oz) as a fertigation or side dress at pinhead square and a second 1-pint application 21 days later.. Apply 8 oz per acre foliar application at pinhead square and second 1-pint application 21 days later. For In-Furrow application, apply 2 oz per acre then follow with a 1/2-pint per acre (8 oz) application at pin square by fertigation, side dress or foliar. Total application should not exceed 2-pints per acre. **DRYLAND COTTON:** Apply 2 oz per acre In-Furrow, OR apply 8 oz per acre at pinhead square and a second 8 oz application 21 days later.

COLE CROPS, such as Broccoli, Broccoli Raab, Brussels Sprouts, Cabbage (Chinese), Cauliflowers, Cavalo Broccoli, Collards, Kale, Kohlrabi, Mizuna, Mustard Greens, Mustard Spinach, Rape Greens: Apply 4 oz per acre as a fertigation or foliar application 2-3 days after transplanting or at 2-3 true leaf stage followed by a second application 10-14 days later. For heading crops such as Broccoli, Cauliflowers, and Brussel Sprouts, apply 4 oz at pre-bloom and repeat application during early head sizing period for 1 additional application.

CUCURBIT VEGETABLES such as Cantaloupe, Chayote, Chinese Waxgourd, Citron Melon, Cucumbers, Gherkin, Edible Gourds, Muskmelon, Pumpkin, Summer Squash, Watermelon, and Winter Squash: Apply 4 oz per acre as a fertigation or foliar application when plants are 3-5 leaf stage or 3-4 days after transplanting. Apply 8 oz per acre at first flower (e.g. first male for cantaloupe). Repeat 14 to 21 days later for last application. Do Not exceed 1 pints per acre over crop cycle.

FRUITING VEGETABLES such as Eggplant, Groundberry, Pepino, Peppers (such as Bell, Chile, Pimento and Sweet Peppers), Tomatillo and Tomatoes: Apply 4 oz per acre as a fertigation or foliar application when plants are 4-6 leaf stage or 3-4 days after transplanting. Apply 4 oz per acre at first flower and again 14 to 21 days later. Do not exceed 16 oz per acre over the crop cycle.

GRAPES, such as Dried, Raisin, Seedless, and Seeded Table Grapes: Apply 8 oz per acre as a fertigation or foliar application, not exceed 1-pint (16 oz) per crop cycle. For Seedless and Seeded Grapes apply after thinning and bloom set has past. Apply second application 14 – 21 days after first application.

GRAPES (Wine): Apply 8 oz per acre as a fertigation or foliar application, followed by a second 8 oz application in 14-21 days. For Seeded Grapes, apply at the start of bloom. Total applications should not exceed 1-pint per acre (16 oz) per crop cycle.

HOPS, all varieties: Apply 12 oz. per acre early growth stage via drip irrigation, followed by 16 oz. per acre med to late growth stage. Apply 16 oz per acre foliar at early lateral development.

LEAFY VEGETABLES such as Amaranth, Arugula, Asparagus, Cardoon, Celery, Celtuce, Chervil, Chrysanthemum, Corn Salad, Cress (garden and upland), Dandelion, Dock, Endive, Fennel, Lettuce (head and leaf), Orach, Parsley, Purslane (garden and winter), Radichio, Rhubarb, Spinach, and Swiss Chard: Apply 4 oz per acre by fertigation or foliar application when plants are at least 3-5 leaves or post transplanting by 3-4 days. NOTE: Applications in excess of 1 pint per acre can cause a delay to harvest.

PINEAPPLES: Apply 1-pint per acre by fertigation or foliar application 18 days after floral induction. Repeat in 30 days. Best performance is achieved with 2-pints over the crop cycle.

POME FRUIT such as Apple, Crabapple, Loquat, Mayhew, Pear (including Oriental), Quince: Apply 8 oz per acre by fertigation or foliar application. Apply foliar with early sizing sprays post bloom. Use as bloom for Crabapple pollinators. Do not exceed a total of 1-pint per acre over the crop cycle.

POTATOES, ROOT AND TUBER VEGETABLES such as Arracha, Arrowroot, Artichoke, Canna (edible), Cassava, Chayote (root), Garlic, Ginger, Onion, Potato, Sweet Potato, Tanier, Yam Bean, Yam Root Vegetables such as Beet (garden & sugar), Burdock (edible), Carrot, Celeriac, Chervil, Chicory, Ginseng, Horseradish, Parsnip, Rutabaga, Salsify, Skirret, and Turnip: Apply 4 oz per acre by fertigation In-Furrow. Apply 12 oz at Hilling by fertigation or foliar (early leaf stages). Maximum of 1.5-pints per acre over the crop cycle.

SOFT FRUIT: Blueberry; Caneberry such as, Raspberry (Red & Black), etc. Apply via drip irrigation or foliar at 8-16 oz per acre starting at pre-bloom followed 2-3 weeks after second application (typically petal fall to early sizing. Depending on varieties a third application at about two weeks after second application or no later than 2 weeks pre-harvest.

STRAWBERRIES: Apply 2 oz. per acre thru fertigation every 7-14 days starting 1-week after transplanting. When flower buds begin apply 2 oz by foliar or drip application every 7-14 days.

STONE FRUIT such as Apricot, Cherry (sweet & tart), Nectarine, Peach, Plum, Plumcot, Prune (dry and fresh types): A total of 3 applications is suggested. Apply 8 oz per acre by fertigation or foliar method at early pink or red bud stage. Apply 4 oz per acre at petal fall. Apply 4 oz per acre final application 2-3 weeks later. Do not exceed 1 pint per acre per season.

SOYBEANS, SUCCULENT, DRIED BEANS AND PEAS such as Beans (Phaseolus and Vigna spp.), Jackbean, Pea (pisum spp.), Pigeon Pea, Sword Bean, and Dried Cultivars of Bean (Lupinus, Phaseolus, Vigna spp.), Broad Bean, Chickpea, Guar, Lablab Bean, Lentil and Pea: Apply 2 oz per acre In-Furrow. Followed by 4 oz per acre application 3-5 Trifoliates stage. OR apply 4 oz per acre at very early vegetative stages such as 3-5 trifoliates or in early reproductive stages followed by 4 oz per acre 14-21 days later. Do not apply more than 8 oz per crop cycle.

TREE NUTS, such as Almond, Beech Nut, Brazil Nut, Butternut, Cashew, Chestnut, Chinquapin, Filbert, Hickory Nut, Macadamia Nut, Pecan, Walnut (Black and English), and Pistachio: Apply 1 (16 oz.) per acre by fertigation or foliar method at the "pink" stage for Almonds. Apply at early flower on other Nut and Pistachio Crops. Apply at a second application of 8 oz per acre after petal fall. Apply a maximum of 2 pints per acre over the crop cycle.

TROPICAL TREE FRUITS such as Avocado, Cherimoya, Coffee, Guava, Lychee, Mango, and Papaya: Apply 8 oz per acre by Foliar method at early pre-flowering to first flower. Repeat 8 oz Foliar application prior to summer nut drop or following a cold snap or hail or excessive rains. Do not apply more than 1-pint per acre over crop cycle.

For Hazardous Materials [or Dangerous Goods] Incident Spill, Leak, Fire, Exposure, or Accident Call CHEMTREC Day or MNight 1-800-424-9300/ +1 703-527-3887 CCN816681.

STORAGE and DISPOSAL

STORAGE & DISPOSAL: Do not contaminate water, food or feed by storage, disposal or cleaning of equipment. Store in a safe manner. Store in original container only and keep tightly sealed when not in use. Dispose of unused product and empty containers in accordance with federal, state and local regulations.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY BEFORE BUYING OR USING THIS PRODUCT, read the Directions for Use and the following Conditions of Sale and Limitation of Warranty and Liability. By buying or using this product, the buyer or user accepts the following Conditions of Sale and Limitation of Warranty and Liability, which no employee or agent of OMEX® AGRIFLUIDS, INC. or the seller is authorized to vary in any way.

Follow the Directions for Use of this product carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop or other plant injury, ineffectiveness, or other unintended consequences may result from such risks as weather or crop conditions, mixture with other chemicals not specifically identified in this product's label, or use of this product contrary to the label instructions, all of which are beyond the control of OMEX® AGRIFLUIDS, INC. and the seller. The buyer or use of this product assumes all such inherent risks.

Subject to the foregoing inherent risks, OMEX® AGRIFLUIDS, INC. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, when the product is used in strict accordance with such Directions for Use under normal conditions of use. EXCEPT AS WARRANTED IN THIS LABEL AND TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THIS PRODUCT IS SOLD "AS IS", AND OMEX™ AGRIFLUIDS, INC. MAKES NO OTHER WARRANTY, EXPRESS OR IMPLIED, INCLUDING BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, OR ELIGIBILITY OF THIS PRODUCT FOR ANY PARTICULAR TRADE USAGE.

IN THE UNLIKELY EVENT THAT BUYER OR USER BELIEVES THAT OMEX® AGRIFLUIDS, INC. HAS BREACHED A WARRANTY CONTAINED IN THIS LABEL AND TO THE EXTENT REQUIRED BY APPLICABLE LAW, BUYER OR USER MUST SEND WRITTEN NOTICE OF ITS CLAIM TO THE FOLLOWING ADDRESS: OMEX® AGRIFLUIDS, INC., ATTENTION LAW DEPARTMENT, PO BOX 815, SELMA, CA 93662.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE BUYER'S OR USER'S EXCLUSIVE REMEDY FOR ANY INJURY, LOSS, OR DAMAGE RESULTING FROM THE HANDLING OR USE OF THIS PRODUCT, INCLUDING BUT NOT LIMITED TO CLAIMS OF BREACH OF WARRANTY OR CONTRACT, NEGLIGENCE, STRICT LIABILITY, OR OTHER TORTS, SHALL BE LIMITED TO ONE OF THE FOLLOWING, AT THE ELECTION OF OMEX® AGRIFLUIDS, INC. OR THE SELLER: DIRECT DAMAGES NOT EXCEEDING THE PURCHASE PRICE OF THE PRODUCT OR REPLACEMENT OF THE PRODUCT TO THE EXTENT CONSISTENT WITH APPLICABLE LAW. OMEX® AGRIFLUIDS, INC. AND THE SELLER SHALL NOT BE LIABLE TO THE BUYER OR USER OF THIS PRODUCT FOR ANY CONSEQUENTIAL, SPECIAL, OR INDIRECT DAMAGES, OR DAMAGES IN THE NATURE OF A PENALTY.

Tests have not been carried out under all possible use conditions and Seller cannot and does not represent or warrant that the goods are compatible with all other chemicals or under all use conditions.

Notice: Neither the manufacturer or the seller make any warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all responsibility for safety and use not in accordance with directions.

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.html>