



Bayer CropScience LP
P.O. Box 12014
2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709
1-866-99BAYER (1-866-992-2937)
<http://www.bayercropscienceus.com>

OSPREY™ Herbicide
EPA Reg No. 264-802

OSPREY™ Herbicide for use on fall-sown or winter wheat from emergence up to the jointing stage of development for partial control of jointed goatgrass and quackgrass.

PRODUCT BULLETIN

For Use and Distribution Only in the State of Montana

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or accepted by the U.S. Environmental Protection Agency. It is a violation of federal law to use this product in a manner inconsistent with its label. This product bulletin must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

Osprey™ Herbicide is labeled for control of wild oat and Persian darnel with partial control of numerous brome species. Light to moderate infestations of jointed goatgrass and quackgrass have exhibited partial control when treated with Osprey™ Herbicide, particularly when tankmixed with EC broadleaf herbicides.

**Partial Control of Jointed Goatgrass and Quackgrass in
Winter Wheat with OSPREY™ Herbicide**

Product	Rate	Crop	Weed
OSPREY™ Herbicide + Methylated Seed Oil	4.75 ounces/acre + 1.3 – 1.5 pints/acre	Apply OSPREY™ Herbicide to winter wheat from emergence up to the jointing stage of wheat.	For partial control of jointed goatgrass (<i>Aegilops cylindrica</i>) and quackgrass (<i>Agropyron repens</i>).

Restrictions and Limitations

- Partial control is defined as stunted growth and a reduction in number as compared to non-treated areas. However, performance may not be commercially acceptable.
- A recommended adjuvant must be tank-mixed with Osprey™ Herbicide according to the guidelines on the federal label.
- Applications should be made to young, actively growing jointed goatgrass and quackgrass. Partial control may be reduced when weeds are under stress due to severe weather conditions, drought, very cold temperatures, etc.
- Do not apply within 30 days of harvesting wheat forage, and 60 days for hay, grain and straw.
- Follow all applicable use directions, weed and crop application timings, precautions and restrictions on the product label for Osprey™ Herbicide.

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577
For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)
or
Visit our worldwide web site at www.bayercropscienceus.com

*As with any crop-protection product, always read and follow label directions.
For additional information call toll-free 1-866-99BAYER (1-866-992-2937).*