

**FIELD & VEGETABLE CROPS**

Make the first application after transplanting, thinning or at second true leaf stage. Apply 1 to 2 subsequent applications at 10 to 15 days intervals as needed.

**Tomatoes & Peppers, Broccoli, Cauliflowers (Brassica), Melons, Cucumbers (Cucurbits):** Use 2 to 4 pints per acre (2.2-4.5 Liters/HA) per application.

**Lettuce, Celery, Spinach, Endives, Carrots, Parsleys & Other Leafy Crops:** Use 2 to 4 pints per acre (2.2-4.5 ltrs./hectare) per application.

**Beans & Peas, Corn:** Use 2 to 4 pints per acre (2.2-4.5 Liters/HA) per application.

**Strawberries:** Use 2 to 4 pints per acre (2.2-4.5 ltrs./hectare) per application. Make the first application after transplanting. Apply at 30-day intervals or as needed to supplement nutritional requirements.

**Potatoes, Onions and Other Tuber Crops:** Improves foliage color as well as tuber development. Use 2 to 4 pints per acre (2.2-4.5 Liters/HA) per application.

**GRAPES, HOPS AND OTHER VINE CROPS**

**Grapes, Hops:** Use 2 to 4 pints per acre (2.2- 4.5 ltrs./hectare) per application.

**Table Grapes:** Make first application at bloom. Apply subsequent applications at bunch preclosing, veraison and 2 to 3 weeks prior to harvest or as needed to supplement nutritional requirements.

**Wine Grapes:** Make first application at approximately 5% bloom. Apply subsequent applications at bunch preclosing, veraison and 2 to 3 weeks prior to harvest or as needed to supplement nutritional requirements.

**Raspberries, Blackberries, Blueberries, Kiwis and Other Cranberries:** Use 2 to 4 pints per acre (2.2-4.5 ltrs. hectare) per application. Make first application at pre-bloom. Make subsequent applications at 30-day intervals up to harvest or as needed to supplement nutritional requirements.

**TREE CROPS: FRUIT, NUT AND CITRUS**

**Almonds, Walnuts and Other Nut Crops:** Use 2 to 4 pints per acre (2.2-4.5 ltrs./hectare) per application. Make the first application at green tip. Make subsequent applications at 30 day intervals up to nut fill or as needed to supplement nutritional requirements.

# PHOS PRO<sup>®</sup>

## ZINC

**0-16-0****GUARANTEED ANALYSIS**

Available Phosphate (P<sub>2</sub>O<sub>5</sub>) ..... 16.00%      Zinc (Zn) ..... 5.50%

Derived from Zinc Phosphite

**PHOS PRO is a unique nutritional supplement that supplies phosphate in a highly concentrated soluble form for foliar or soil applications. Properly timed applications will help result in beneficial plant growth, improved crop quality and yield increase.**

Upon foliar application, the phosphite ions (PHOS PRO) are taken up directly by the plant foliage and may undergo a degree of conversion to phosphate ions or will be used directly by the plants as phosphite ions. Phosphorous acid products (PHOS PRO) are for use as a supplemental fertilizer treatment.

**PHOS PRO Zinc provides advantageous buffering in the preferred range of pH.**

Water pH	0.50% PHOS PRO Zinc Solution
Start	Change
pH 7.86	6.10

**DIRECTIONS**

**Aerial Application:** Use 1 to 4 pints per acre in minimum of 10 gals of water (1.1 to 4.5 ltrs. per hectare/95 ltrs.).

**Ground Application:** Use 1 to 4 pints per acre in minimum of 20 gals of water (1.1 to 3.2 ltrs. per hectare). Do not exceed 2% volume/volume product concentration.

i.e.: 1 pint/acre ..... use min. 6.25 gals. of water/acre

1.1 liters/hectare ..... use min. 56.7 liters of water/hectare

**Soil Application:** Use 3 to 4 pints per acre (3.4 to 4.5 liters per hectare), applied through low water volume irrigation systems or as a side dressing.

**Compatibility:** PHOS PRO is compatible with most commonly used pesticides and nutrient products. However, prior to using previously untried mixes, perform a jar test to evaluate potential product incompatibilities.

**NET CONTENT: 2.5 GALLONS (9.46 L)****NET WEIGHT: 24.95 LBS. (11.3 KG)**

**Plums, Nectarines and Other Stone Fruits:** Use 2 to 4 pints per acre (2.2-4.5 ltrs./hectare) per application. Make the first application at green tip or pink bud. Make subsequent applications at fruit set or jacket fall, fruit thinning and 2 to 3 weeks prior to harvest or as needed to supplement nutritional requirements.

**Apples, Pears and Other Pome Fruits:** Use 2 to 4 pints per acre (2.2-4.5 ltrs./hectare) per application. Make the first application at green tip. Make subsequent applications at 30-day intervals up to harvest or as needed to supplement nutritional requirements.

**Citrus and Avocados:** Use 2 to 4 pints per acre (2.2-4.5 ltrs./hectare) per application. Make the first application at pre-bloom. Make subsequent applications at mid crop and 3 to 4 weeks prior to harvest or as needed to supplement nutritional requirements.

**Olives:** Use 2 to 4 pints per acre (2.2-4.5 ltrs./hectare) per application. Make the first application at pre-bloom. Apply subsequent applications at 30-day intervals to harvest or as needed to supplement nutritional requirements.

**CAUTION:**

**ACIDIC MIXTURE MAY BE IRRITATING TO EYES AND TO SKIN.**

**KEEP OUT OF REACH OF CHILDREN**

Avoid breathing and prolonged exposure to dermis. Do not apply when weather conditions favor drift from the areas treated. Do not apply when runoff is likely to occur. Do not use in a manner or at a time other than in accordance with label directions because animal, plant or crop injury, illegal residue or other results may occur.

Information regarding the contents and levels of metals in this product is available on the internet at: <http://www.aapfco.org/metals.htm>

Manufactured By

**GROW  MORE<sup>®</sup>**

Grow More, Inc.  
15600 New Century Drive  
Gardena, CA 90248  
[www.growmore.com](http://www.growmore.com)

