

# BOOSTER 42 FOLIAR SPRAY 8-30-4

## **GUARANTEED ANALYSIS:**

TOTAL NITROGEN (N)	8.0%
8.0% Ammoniacal Nitrogen	
AVAILABLE PHOSPHATE (P <sub>2</sub> O <sub>5</sub> )	.30.0%
SOLUBLE POTASH (K <sub>2</sub> O) <sup>2</sup> 37	4.00%
Iron (Fe)	0.10%
Zinc (Zn)	

Derived from Ammonium Polyphosphate, Potassium Polyphosphate, Ferrous Sulfate and Zinc Sulfate.

Weight per gallon 11.4 lbs. per gallon @ 68°F.



# KEEP OUT OF REACH OF CHILDREN WARNING

CAUSES SKIN IRRITATION. CAUSES EYE IRRITATION. MAY CAUSE RESPIRATORY IRRITATION.

**PRECAUTIONARY STATEMENTS:** Avoid breathing fumes / gas / mist / vapors / spray. Wash thoroughly after handling. Use only outdoors or in a well-ventilated area. Call a poison center / doctor if you feel unwell. Specific treatment see First Aid section on this label.

**PERSONAL PROTECTIVE EQUIPMENT:** Wear protective gloves / protective clothing / eye protection / face protection. Take off contaminated clothing and wash before reuse.

FIRST AID: IF ON SKIN Wash with plenty of water. If skin irritation occurs get medical advice / attention. IF INHALED Remove person to fresh air and keep comfortable for breathing. IF IN EYES Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists get medical advice / attention.

STORAGE AND DISPOSAL: STORAGE Store in a well-ventilated place. Keep container tightly closed. Store locked up. DISPOSAL Dispose of contents / container in accordance with local / regional / national regulations.

# **DIRECTIONS FOR USE**

**PHT BOOSTER 42** is a liquid plant food recommended for use as a foliar spray during periods of rapid growth. It may be applied by air or by ground. Dilute **PHT BOOSTER** 42 in 5-10 gallons of water for air application and in 20-200 gallons for ground rigs.

Apply when there is ample foliage for nutrient absorption. Best results are obtained when sprayed during late evening or early morning -- not in the heat of the day.

A wetting agent is recommended for thorough coverage. SHAKE WELL BEFORE USING.

#### **RECOMMENDATIONS FOR USE**

**GRAPES:** Apply 1-3 quarts per acre prior to bloom. Repeat applications as needed up to six weeks before harvest.

**TOMATOES:** Apply 2-3 quarts per acre prior to first bloom. Repeat at 2-3 week intervals.

**TREE CROPS:** Apply one gallon per acre at bud stage prior to bloom. Repeat after petal fall is complete and again in 4-6 weeks.

**VEGETABLES, FIELD CROPS:** Apply 2-3 quarts per acre prior to bloom. Repeat at 10-30 day intervals as needed.

DISCLAIMER OF WARRANTIES: Seller warrants that the chemical composition of this product conforms to the chemical description given on this label. THIS WARRANTY IS EXPRESSLY IN LIEU OF ALL OTHER WARRANTIES AND REPRESENTATIONS EXPRESSED, IMPLIED, OR STATUTORY, INCLUDING WARRANTIES OF MERCHANTABILITY AND FITNESS FOR A PARTICULAR USE. Timing, rate and method of application, weather and crop conditions, mixtures with chemicals not specifically recommended on this label or an accompanying written recommendation are beyond the control of seller. Buyer assumes all risks of use, storage and handling of this material not in strict accordance with directions given herewith. Buyer further agrees in the event of damages arising from the use of this product to accept a replacement of the product or a refund of the purchase price of the product, at buyer's option, as full discharge of seller's liability. No one is authorized to make any other warranty, guarantee or directions concerning this product, and no such warranties, guarantees or directions shall be valid or binding upon seller.

GHS 4-3-2015

**NET CONTENTS: 5 GALLONS (18.9 L)** 



## J.R. SIMPLOT COMPANY

P.O. Box 70013, Boise, ID 83707-0113 (208)336-2110 GUARANTOR:

P.O. Box 198, Lathrop, CA 95330-0198