



# Folo Spray 6-30-30

**WITH  
MICRONUTRIENTS****GUARANTEED ANALYSIS:**

TOTAL NITROGEN (N) .....	6.00%
2.0% Ammoniacal Nitrogen	
4.0% Nitrate Nitrogen	
AVAILABLE PHOSPHATE (P <sub>2</sub> O <sub>5</sub> ) .....	30.00%
SOLUBLE POTASH (K <sub>2</sub> O) .....	30.00%
Boron (B) .....	0.02%
Copper (Cu) .....	0.05%
0.05% Chelated Copper	
Iron (Fe) .....	0.10%
0.10% Chelated Iron	
Manganese (Mn) .....	0.05%
0.05% Chelated Manganese	
Molybdenum (Mo) .....	0.0005%
0.0005% Water Soluble Molybdenum	
Zinc (Zn) .....	0.05%
0.05% Chelated Zinc	

Derived from Monoammonium Phosphate, Monopotassium Phosphate, Potassium Nitrate, Sodium Borate, Copper EDTA, Iron EDTA, Manganese EDTA, Sodium Molybdate, and Zinc EDTA.

**KEEP OUT OF REACH OF CHILDREN****WARNING**

CAUSES EYE IRRITATION. MAY INTENSIFY FIRE; OXIDIZER

**PRECAUTIONARY STATEMENTS:** Wash hands thoroughly after handling. Keep away from heat. Keep/Store away from clothing, combustible materials. Take any precaution to avoid mixing with combustible materials.

**PERSONAL PROTECTIVE EQUIPMENT:** Wear protective gloves / eye protection / face protection.

**FIRST AID: IF IN EYES** Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists: Get medical advice / attention.

**IN CASE OF FIRE:** Use chemical type foam, carbon dioxide, dry chemical, water fog or spray to extinguish.

**DISPOSAL** Dispose of contents / container in accordance with Federal, state, and local regulations.

**GENERAL USE DIRECTIONS**

**DILUTION RATE:** Unless otherwise specified, use the rate per acre and dilute as follows:

**AIR:** 5-30 gallons of water per acre.

**DILUTE:** 100 plus gallons of water per acre.

**CONCENTRATE:** 10-100 gallons of water per acre.

Do not exceed 2 lbs. of product per gallon of water. For backpack sprayers use 4 teaspoons per gallon of water.

When temperatures are cold allow extra time for this product to completely dissolve. Do not use when temperatures are expected to exceed 90°F or when plants are under stress conditions such as heat or water stress.

THIS PRODUCT WILL LOWER THE pH OF THE SPRAY SOLUTION.

**DO NOT EXCEED RECOMMENDED AMOUNTS PER ACRE.**

**MIXING DIRECTIONS:** Add approximately ½ water to tank before adding FOLO SPRAY 6-30-30, agitate thoroughly while adding remaining water, add pesticide last. Water temperature plays an important role in determining fertilizer solubility. The higher the water temperature the greater the solubility and quicker the dissolving time. As fertilizers are added to the solution, there is an immediate decrease in water temperature. It is important when preparing concentrated solutions to allow for this temperature drop to ensure that all the fertilizer is properly dissolved.

**COMPATIBILITY:** FOLO SPRAY 6-30-30 is compatible with most commonly used pesticides. However, a compatibility test is recommended before full scale mixing, if the desired combination has not been used in the past. Do not combine with copper fungicide or other strongly alkaline materials.

**RECOMMENDATIONS FOR USE**

**TREE FRUIT, NUT and VINE CROPS such as:** Apples, Almonds, Apricots, Avocados, Citrus (all kinds), Cherries, Filberts, Peaches, Pears, Plums, Prunes, Pecans, Pistachios, Nectarines, Walnuts, Tropical and Sub-Tropical Fruits, Bushberries, Grapes and other vine crops. Apply early in the growing season and continue throughout the season to supplement normal fertility programs.

Apply as a pre-bloom and post-bloom spray. Rate per acre will depend on the amount of carrier applied per acre. In dilute spray applications, use 2 to 5 lbs. per 100 gallons of water. In concentrated spray applications, use 5 to 15 lbs. per acre. Avoid combining with materials that may cause russetting on fruit. A post harvest application of 10 to 40 lbs. per acre can enhance tree vigor and increase nutrient levels within fruiting spurs for the following season. **CAUTION:** Do not use FOLO SPRAY 6-30-30 in late season sprays where fruit color and maturity is delayed by nitrogen applications.

**FIELD, VEGETABLE and HERB CROPS:** As a foliar spray use 5 to 10 lbs. per acre. Applications may be made at 7 to 10 day intervals depending on the amount of growth desired or when supplemental fertilization is required. Depending on the crop, the first treatment should be made when the plants are 3 to 4 weeks old or when there is sufficient foliage for spraying. These rates may be applied to field crops such as: Cotton, Soybeans, Peanuts, Corn, Alfalfa, Sorghum, and forage crops or to vegetable crops such as: Tomatoes, Peppers, Cucumbers, Beans, Lettuce, Celery, Melons, Squash, Radishes, Onions, Broccoli, Cabbage, Spinach, Cauliflower, Strawberries, and Potatoes.

**COMMERCIAL NURSERY and GREENHOUSE:** As a direct foliar spray use 2 to 4 lbs. per 100 gallons of water or 4 teaspoons per gallon of water.

**Greenhouse Cut Flowers, Flats, Small Pots:** Use 2 lbs. per 100 gallons of water for intermittent feeding (every 2 weeks). For constant feeding (every watering) use 4 to 8 oz. per 100 gallons of water.

**Container Foliage Plants:** Use up to 3 1/2 lbs. per 100 gallons of water for intermittent feeding. For constant feeding use 8 to 12 oz. per 100 gallons of water.

**Orchids:** For periodic feeding use 5 to 6 lbs. in 100 gallons of water. For constant feeding use 1 to 1 1/2 lbs. per 100 gallons of water. Starter and Transplant: Use at the rate of 1 lb. per 25 gallons of water as plants are being set out in the field. Apply sufficient solution for proper soil moisture.

**LANDSCAPE TREES AND SHRUBS, TURF:** For foliar spray or root feeding use 2 lbs. per 100 gallons of water. For seeding new turf, 1/2 lb. may be mixed with each pound of grass seed. Apply by spreader or broadcast evenly over 200 sq. ft.

**LAWNS, AND GOLF GREENS:** While soil is wet apply 1 lb. per 2,000 sq. ft. dissolved in 10 or more gallons of water. Color and rate of growth will determine frequency of application.

Warning: This product contains a chemical known to the State of California to cause cancer, birth defects or other reproductive harm. Proposition 65, the Safe Drinking Water and Toxic Enforcement Act of 1986, requires notification of potential exposure to substances identified by the State of California as causing cancer, birth defects or other reproductive harm.

Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.regulatory-info-jr.com>

**DISCLAIMER OF WARRANTIES:** Seller warrants that the chemical composition of this product conforms to the chemical description given on this label. THIS WARRANTY IS EXPRESSLY IN LIEU OF ALL OTHER WARRANTIES AND REPRESENTATIONS EXPRESSED, IMPLIED, OR STATUTORY, INCLUDING WARRANTIES OF MERCHANTABILITY AND FITNESS FOR A PARTICULAR USE. Timing, rate and method of application, weather and crop conditions, mixtures with chemicals not specifically recommended on this label or an accompanying written recommendation, are beyond the control of seller. Buyer assumes all risks of use, storage and handling of this material not in strict accordance with directions given herewith. Buyer further agrees in the event of damages arising from the use of this product to accept a replacement of the product or a refund of the purchase price of the product, at buyer's option, as full discharge of seller's liability. No one is authorized to make any other warranty, guarantee or directions concerning this product, and no such warranties, guarantees or directions shall be valid or binding upon seller.

GHS\_R2-15-15

**NET WEIGHT: 25 POUNDS (11.34 Kg.)****J. R. SIMPLOT COMPANY**

P.O. Box 70013, Boise, ID 83707-0113 • 208-336-2110  
GUARANTOR • P.O. Box 198, Lathrop, CA 95330